

## Product datasheet for **TL314418V**

### **BTN1A1 Human shRNA Lentiviral Particle (Locus ID 696)**

#### **Product data:**

|                      |   |
|----------------------|---|
| <b>Product Type:</b> | shRNA Lentiviral Particles  |
| <b>Product Name:</b> | BTN1A1 Human shRNA Lentiviral Particle (Locus ID 696)   |
| <b>Locus ID:</b>     | 696   |
| <b>Synonyms:</b>     | BT; BTN; BTN1   |
| <b>Vector:</b>       | pGFP-C-shLenti (TR30023)  |
| <b>Format:</b>       | Lentiviral particles  |
| <b>Components:</b>   | BTN1A1 - Human shRNA lentiviral particles (4 unique 29mer target-specific shRNA, 1 scramble control), 0.5 ml each, >10 <sup>7</sup> TU/ml.  |
| <b>RefSeq:</b>       | <a href="#">NM_001732</a> , <a href="#">NM_001732.1</a> , <a href="#">NM_001732.2</a> , <a href="#">BC096312</a> , <a href="#">BC096313</a> , <a href="#">BC096314</a> , <a href="#">BC096315</a>   |
| <b>UniProt ID:</b>   | <a href="#">Q13410</a>  |
| <b>Summary:</b>      | Butyrophilin is the major protein associated with fat droplets in the milk. It is a member of the immunoglobulin superfamily. It may have a cell surface receptor function. The human butyrophilin gene is localized in the major histocompatibility complex (MHC) class I region of 6p and may have arisen relatively recently in evolution by the shuffling of exons between 2 ancestral gene families [provided by RefSeq, Jul 2008] |
| <b>shRNA Design:</b> | These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact <a href="mailto:techsupport@origene.com">techsupport@origene.com</a> . If you need a special design or shRNA sequence, please utilize our <a href="#">custom shRNA service</a> .  |



[View online »](#)

**Performance  
Guaranteed:**

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at [techsupport@origene.com](mailto:techsupport@origene.com). Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).