

Product datasheet for TL314429V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD147 (BSG) Human shRNA Lentiviral Particle (Locus ID 682)

Product data:

Product Type: shRNA Lentiviral Particles

Product Name: CD147 (BSG) Human shRNA Lentiviral Particle (Locus ID 682)

Locus ID: 682

Synonyms: 5F7; CD147; EMMPRIN; EMPRIN; HAb18G; OK; SLC7A11; TCSF

Vector: pGFP-C-shLenti (TR30023)

Format: Lentiviral particles

Components: BSG - Human shRNA lentiviral particles (4 unique 29mer target-specific shRNA, 1 scramble

control), 0.5 ml each, >10^7 TU/ml.

RefSeq: NM 001322243, NM 001728, NM 198589, NM 198590, NM 198591, NM 198589.1,

NM 198589.2, NM 198591.1, NM 198591.2, NM 001728.1, NM 001728.2, NM 001728.3,

NM 198590.1, NM 198590.2, BC009040, BC009040.2, NM 198590.3, NM 001728.4,

NM 198591.3, NM 198589.3

UniProt ID: P35613

Summary: The protein encoded by this gene, basigin, is a plasma membrane protein that is important in

spermatogenesis, embryo implantation, neural network formation, and tumor progression. Basigin is also a member of the immunoglobulin superfamily, ubiquitously expressed in various tissues. Multiple transcript variants encoding different isoforms have been found for

this gene. [provided by RefSeq, May 2020]

shRNA Design: These shRNA constructs were designed against multiple splice variants at this gene locus. To

be certain that your variant of interest is targeted, please contact <u>techsupport@origene.com</u>. If you need a special design or shRNA sequence, please utilize our <u>custom shRNA service</u>.






Performance Guaranteed:

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.


For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).

Product images:




GFP signal was observed under microscope at 48 hours after transduction of TL314429A virus into HEK293 cells. TL314429A virus was prepared using lenti-shRNA TL314429A and [TR30037] packaging kit.






GFP signal was observed under microscope at 48 hours after transduction of TL314429A virus into HEK293 cells. TL314429A virus was prepared using lenti-shRNA TL314429A and [TR30037] packaging kit.




GFP signal was observed under microscope at 48 hours after transduction of TL314429B virus into HEK293 cells. TL314429B virus was prepared using lenti-shRNA TL314429B and [TR30037] packaging kit.




GFP signal was observed under microscope at 48 hours after transduction of TL314429B virus into HEK293 cells. TL314429B virus was prepared using lenti-shRNA TL314429B and [TR30037] packaging kit.






GFP signal was observed under microscope at 48 hours after transduction of [TL314429C] virus into HEK293 cells. [TL314429C] virus was prepared using lenti-shRNA [TL314429C] and [TR30037] packaging kit.




GFP signal was observed under microscope at 48 hours after transduction of [TL314429C] virus into HEK293 cells. [TL314429C] virus was prepared using lenti-shRNA [TL314429C] and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of [TL314429D] virus into HEK293 cells. [TL314429D] virus was prepared using lenti-shRNA [TL314429D] and [TR30037] packaging kit.





GFP signal was observed under microscope at 48 hours after transduction of [TL314429D] virus into HEK293 cells. [TL314429D] virus was prepared using lenti-shRNA [TL314429D] and [TR30037] packaging kit.