

Product datasheet for **TL315477V**

BIRC5 Human shRNA Lentiviral Particle (Locus ID 332)

Product data:

Product Type:	shRNA Lentiviral Particles
Product Name:	BIRC5 Human shRNA Lentiviral Particle (Locus ID 332)
Locus ID:	332
Synonyms:	API4; EPR-1
Vector:	pGFP-C-shLenti (TR30023)
Format:	Lentiviral particles
Components:	BIRC5/Survivin ^Δ - Human shRNA lentiviral particles (4 unique 29mer target-specific shRNA, 1 scramble control), 0.5 ml each, >10 ⁷ TU/ml.
RefSeq:	<u>NM_001012270</u> , <u>NM_001012271</u> , <u>NM_001168</u> , <u>NM_001168.1</u> , <u>NM_001168.2</u> , <u>NM_001012270.1</u> , <u>NM_001012271.1</u> , <u>BC034148</u> , <u>BC034148.1</u> , <u>BC000784</u> , <u>BC008718</u> , <u>BC065497</u> , <u>BM476145</u> , <u>NM_001012271.2</u> , <u>NM_001012270.2</u>
UniProt ID:	<u>O15392</u>
Summary:	This gene is a member of the inhibitor of apoptosis (IAP) gene family, which encode negative regulatory proteins that prevent apoptotic cell death. IAP family members usually contain multiple baculovirus IAP repeat (BIR) domains, but this gene encodes proteins with only a single BIR domain. The encoded proteins also lack a C-terminus RING finger domain. Gene expression is high during fetal development and in most tumors, yet low in adult tissues. Alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, Jun 2011]
shRNA Design:	These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact techsupport@origene.com . If you need a special design or shRNA sequence, please utilize our custom shRNA service .




[View online »](#)


**Performance
Guaranteed:**

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.


For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).

Product images:


GFP signal was observed under microscope at 48 hours after transduction of TL315477A virus into HEK293 cells. TL315477A virus was prepared using lenti-shRNA TL315477A and [TR30037] packaging kit.




GFP signal was observed under microscope at 48 hours after transduction of TL315477A virus into HEK293 cells. TL315477A virus was prepared using lenti-shRNA TL315477A and [TR30037] packaging kit.




GFP signal was observed under microscope at 48 hours after transduction of TL315477B virus into HEK293 cells. TL315477B virus was prepared using lenti-shRNA TL315477B and [TR30037] packaging kit.




GFP signal was observed under microscope at 48 hours after transduction of TL315477B virus into HEK293 cells. TL315477B virus was prepared using lenti-shRNA TL315477B and [TR30037] packaging kit.




GFP signal was observed under microscope at 48 hours after transduction of [TL315477C] virus into HEK293 cells. [TL315477C] virus was prepared using lenti-shRNA [TL315477C] and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of [TL315477C] virus into HEK293 cells. [TL315477C] virus was prepared using lenti-shRNA [TL315477C] and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of [TL315477D] virus into HEK293 cells. [TL315477D] virus was prepared using lenti-shRNA [TL315477D] and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of [TL315477D] virus into HEK293 cells. [TL315477D] virus was prepared using lenti-shRNA [TL315477D] and [TR30037] packaging kit.