

Product datasheet for TL316756V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

mtTFA (TFAM) Human shRNA Lentiviral Particle (Locus ID 7019)

Product data:

Product Type: shRNA Lentiviral Particles

Product Name: mtTFA (TFAM) Human shRNA Lentiviral Particle (Locus ID 7019)

Locus ID: 7019

Synonyms: MTDPS15; MTTF1; MTTFA; TCF6; TCF6L1; TCF6L2; TCF6L3

Vector: pGFP-C-shLenti (TR30023)

Format: Lentiviral particles

Components: TFAM - Human shRNA lentiviral particles (4 unique 29mer target-specific shRNA, 1 scramble

control), 0.5 ml each, >10^7 TU/ml.

RefSeq: NM 001270782, NM 003201, NM 012251, NR 073073, NM 003201.1, NM 003201.2,

NM 001270782.1, NM 012251.1, BC126366, BC018628, BC029815, BC104482, BM455062,

NM 003201.3

UniProt ID: Q00059

Summary: This gene encodes a key mitochondrial transcription factor containing two high mobility

group motifs. The encoded protein also functions in mitochondrial DNA replication and repair. Sequence polymorphisms in this gene are associated with Alzheimer's and Parkinson's diseases. There are pseudogenes for this gene on chromosomes 6, 7, and 11. Alternative

splicing results in multiple transcript variants. [provided by RefSeq, Aug 2012]

shRNA Design: These shRNA constructs were designed against multiple splice variants at this gene locus. To

be certain that your variant of interest is targeted, please contact <u>techsupport@origene.com</u>. If you need a special design or shRNA sequence, please utilize our <u>custom shRNA service</u>.




Performance Guaranteed:

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.


For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).

Product images:




GFP signal was observed under microscope at 48 hours after transduction of TL316756A virus into HEK293 cells. TL316756A virus was prepared using lenti-shRNA TL316756A and [TR30037] packaging kit.






GFP signal was observed under microscope at 48 hours after transduction of TL316756A virus into HEK293 cells. TL316756A virus was prepared using lenti-shRNA TL316756A and [TR30037] packaging kit.




GFP signal was observed under microscope at 48 hours after transduction of TL316756B virus into HEK293 cells. TL316756B virus was prepared using lenti-shRNA TL316756B and [TR30037] packaging kit.




GFP signal was observed under microscope at 48 hours after transduction of TL316756B virus into HEK293 cells. TL316756B virus was prepared using lenti-shRNA TL316756B and [TR30037] packaging kit.






GFP signal was observed under microscope at 48 hours after transduction of [TL316756C] virus into HEK293 cells. [TL316756C] virus was prepared using lenti-shRNA [TL316756C] and [TR30037] packaging kit.




GFP signal was observed under microscope at 48 hours after transduction of [TL316756C] virus into HEK293 cells. [TL316756C] virus was prepared using lenti-shRNA [TL316756C] and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of [TL316756D] virus into HEK293 cells. [TL316756D] virus was prepared using lenti-shRNA [TL316756D] and [TR30037] packaging kit.





GFP signal was observed under microscope at 48 hours after transduction of [TL316756D] virus into HEK293 cells. [TL316756D] virus was prepared using lenti-shRNA [TL316756D] and [TR30037] packaging kit.