

Product datasheet for TL318197

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

H19 Human shRNA Plasmid Kit (Locus ID 283120)

Product data:

Product Type: shRNA Plasmids

Product Name: H19 Human shRNA Plasmid Kit (Locus ID 283120)

Locus ID: 283120

Synonyms: ASM; ASM1; BWS; D11S813E; LINC00008; MIR675HG; NCRNA00008; WT2

Vector: pGFP-C-shLenti (TR30023) **E. coli Selection:** Chloramphenicol (34 ug/ml)

Mammalian Cell

Selection:

Puromycin

Format: Lentiviral plasmids

Components: H19 - Human, 4 unique 29mer shRNA constructs in lentiviral GFP vector(Gene ID = 283120).

5µg purified plasmid DNA per construct

29-mer scrambled shRNA cassette in pGFP-C-shLenti Vector, TR30021, included for free.

RefSeq: NR 002196, NR 131223, NR 131224, BC006831, BC007513, BC009853, BC010185, BC013067,

BC015952, BC023213, BC040073, BC040852, BC053636, BC053637, BC063564, BC063626,

BC069247, BC098439, BC106078, BC110657, BC115700

Summary: This gene is located in an imprinted region of chromosome 11 near the insulin-like growth

factor 2 (IGF2) gene. This gene is only expressed from the maternally-inherited chromosome, whereas IGF2 is only expressed from the paternally-inherited chromosome. The product of this gene is a long non-coding RNA which functions as a tumor suppressor. Mutations in this gene have been associated with Beckwith-Wiedemann Syndrome and Wilms tumorigenesis. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Apr 2015]

shRNA Design: These shRNA constructs were designed against multiple splice variants at this gene locus. To

be certain that your variant of interest is targeted, please contact <u>techsupport@origene.com</u>. If you need a special design or shRNA sequence, please utilize our <u>custom shRNA service</u>.






Performance Guaranteed:

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.


For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).

Product images:




GFP signal was observed under microscope at 48 hours after transduction of TL318197A virus into HEK293 cells. TL318197A virus was prepared using lenti-shRNA TL318197A and [TR30037] packaging kit.






GFP signal was observed under microscope at 48 hours after transduction of TL318197A virus into HEK293 cells. TL318197A virus was prepared using lenti-shRNA TL318197A and [TR30037] packaging kit.




GFP signal was observed under microscope at 48 hours after transduction of TL318197B virus into HEK293 cells. TL318197B virus was prepared using lenti-shRNA TL318197B and [TR30037] packaging kit.




GFP signal was observed under microscope at 48 hours after transduction of TL318197B virus into HEK293 cells. TL318197B virus was prepared using lenti-shRNA TL318197B and [TR30037] packaging kit.






GFP signal was observed under microscope at 48 hours after transduction of [TL318197C] virus into HEK293 cells. [TL318197C] virus was prepared using lenti-shRNA [TL318197C] and [TR30037] packaging kit.




GFP signal was observed under microscope at 48 hours after transduction of [TL318197C] virus into HEK293 cells. [TL318197C] virus was prepared using lenti-shRNA [TL318197C] and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of [TL318197D] virus into HEK293 cells. [TL318197D] virus was prepared using lenti-shRNA [TL318197D] and [TR30037] packaging kit.





GFP signal was observed under microscope at 48 hours after transduction of [TL318197D] virus into HEK293 cells. [TL318197D] virus was prepared using lenti-shRNA [TL318197D] and [TR30037] packaging kit.