

## Product datasheet for **TL320459**

### PPAR gamma (PPARG) Human shRNA Plasmid Kit (Locus ID 5468)

#### Product data:

Product Type:	shRNA Plasmids
Product Name:	PPAR gamma (PPARG) Human shRNA Plasmid Kit (Locus ID 5468)
Locus ID:	5468
Synonyms:	CIMT1; GLM1; NR1C3; PPARG1; PPARG2; PPARG5; PPARGgamma
Vector:	pGFP-C-shLenti (TR30023)
E. coli Selection:	Chloramphenicol (34 ug/ml)
Mammalian Cell Selection:	Puromycin
Format:	Lentiviral plasmids
Components:	PPARG - Human, 4 unique 29mer shRNA constructs in lentiviral GFP vector(Gene ID = 5468). 5µg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pGFP-C-shLenti Vector, TR30021, included for free.
RefSeq:	<a href="#">NM_001330615</a> , <a href="#">NM_005037</a> , <a href="#">NM_015869</a> , <a href="#">NM_138711</a> , <a href="#">NM_138712</a> , <a href="#">NM_001354666</a> , <a href="#">NM_001354667</a> , <a href="#">NM_001354668</a> , <a href="#">NM_001354669</a> , <a href="#">NM_001354670</a> , <a href="#">NM_138712.1</a> , <a href="#">NM_138712.2</a> , <a href="#">NM_138712.3</a> , <a href="#">NM_138711.1</a> , <a href="#">NM_138711.2</a> , <a href="#">NM_138711.3</a> , <a href="#">NM_005037.1</a> , <a href="#">NM_005037.2</a> , <a href="#">NM_005037.3</a> , <a href="#">NM_005037.4</a> , <a href="#">NM_005037.5</a> , <a href="#">NM_015869.1</a> , <a href="#">NM_015869.2</a> , <a href="#">NM_015869.3</a> , <a href="#">NM_015869.4</a> , <a href="#">BC006811</a> , <a href="#">BM923992</a> , <a href="#">NM_138712.4</a> , <a href="#">NM_015869.5</a> , <a href="#">NM_138711.4</a> , <a href="#">NM_005037.6</a>
UniProt ID:	<a href="#">P37231</a>
Summary:	This gene encodes a member of the peroxisome proliferator-activated receptor (PPAR) subfamily of nuclear receptors. PPARs form heterodimers with retinoid X receptors (RXRs) and these heterodimers regulate transcription of various genes. Three subtypes of PPARs are known: PPAR-alpha, PPAR-delta, and PPAR-gamma. The protein encoded by this gene is PPAR-gamma and is a regulator of adipocyte differentiation. Additionally, PPAR-gamma has been implicated in the pathology of numerous diseases including obesity, diabetes, atherosclerosis and cancer. Alternatively spliced transcript variants that encode different isoforms have been described. [provided by RefSeq, Jul 2008]
shRNA Design:	These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact <a href="mailto:techsupport@origene.com">techsupport@origene.com</a> . If you need a special design or shRNA sequence, please utilize our <a href="#">custom shRNA service</a> .




[View online »](#)


**Performance  
Guaranteed:**

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.


For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at [techsupport@origene.com](mailto:techsupport@origene.com). Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).

**Product images:**


GFP signal was observed under microscope at 48 hours after transduction of TL320459A virus into HEK293 cells. TL320459A virus was prepared using lenti-shRNA TL320459A and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of TL320459B virus into HEK293 cells. TL320459B virus was prepared using lenti-shRNA TL320459B and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of [TL320459C] virus into HEK293 cells. [TL320459C] virus was prepared using lenti-shRNA [TL320459C] and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of [TL320459D] virus into HEK293 cells. [TL320459D] virus was prepared using lenti-shRNA [TL320459D] and [TR30037] packaging kit.