

Product datasheet for **TL500528**

Dio2 Mouse shRNA Plasmid (Locus ID 13371)

Product data:

Product Type:	shRNA Plasmids
Product Name:	Dio2 Mouse shRNA Plasmid (Locus ID 13371)
Locus ID:	13371
Synonyms:	5DII; AI324267; DIOII
Vector:	pGFP-C-shLenti (TR30023)
E. coli Selection:	Chloramphenicol (34 ug/ml)
Mammalian Cell Selection:	Puromycin
Format:	Lentiviral plasmids
Components:	Dio2 - Mouse, 4 unique 29mer shRNA constructs in lentiviral GFP vector(Gene ID = 13371). 5µg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pGFP-C-shLenti Vector, TR30021, included for free.
RefSeq:	BC125383 , BC125385 , NM_010050 , NM_010050.1 , NM_010050.2 , NM_010050.3
UniProt ID:	Q9Z1Y9



[View online »](#)

Summary:	<p>The protein encoded by this gene belongs to the iodothyronine deiodinase family. It catalyzes the conversion of prohormone thyroxine (3,5,3',5'-tetraiodothyronine, T4) to the bioactive thyroid hormone (3,5,3'-triiodothyronine, T3) by outer ring 5'-deiodination. This gene is highly expressed in brain, placenta and mammary gland. It is thought to be responsible for the 'local' production of T3, and thus important in influencing thyroid hormone action in these tissues. Knockout studies in mice suggest that this gene may play an important role in brown adipose tissue lipogenesis, auditory function, and bone formation. This protein is a selenoprotein containing the non-standard amino acid, selenocysteine (Sec), which is encoded by the UGA codon that normally signals translation termination. The 3' UTRs of selenoprotein mRNAs contain a conserved stem-loop structure, designated the Sec insertion sequence (SECIS) element, that is necessary for the recognition of UGA as a Sec codon, rather than as a stop signal. Unlike the other two members (DIO1 and DIO3) of this enzyme family, the mRNA for this gene contains an additional in-frame UGA codon that has been reported (in human) to function either as a Sec or a stop codon, resulting in two potential isoforms with one or two Sec residues; however, only the upstream Sec (conserved with the single Sec residue found at the active site in DIO1 and DIO3) was shown to be essential for enzyme activity (PMID:10403186). In addition, the lack of conservation of the protein extension past the second TGA codon suggests that the one-Sec containing isoform represents the canonical form. [provided by RefSeq, Oct 2018]</p>
shRNA Design:	<p>These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact techsupport@origene.com. If you need a special design or shRNA sequence, please utilize our custom shRNA service.</p>
Performance Guaranteed:	<p>OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.</p> <p>For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).</p>