

Product datasheet for TL501807

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Ptprj Mouse shRNA Plasmid (Locus ID 19271)

Product data:

Product Type: shRNA Plasmids

Product Name: Ptprj Mouse shRNA Plasmid (Locus ID 19271)

Locus ID: 19271

Synonyms: Al450271; BET; Byp; CD148; DEP-1; Ptpb2; PTPbeta2; RPTPJ; Scc-1; Scc1

Vector: pGFP-C-shLenti (TR30023)

E. coli Selection: Chloramphenicol (34 ug/ml)

Mammalian Cell

Selection:

Puromycin

Format: Lentiviral plasmids

Components: Ptprj - Mouse, 4 unique 29mer shRNA constructs in lentiviral GFP vector(Gene ID = 19271).

5µg purified plasmid DNA per construct

29-mer scrambled shRNA cassette in pGFP-C-shLenti Vector, TR30021, included for free.

RefSeq: NM 001135657, NM 008982, NM 001135657.1, NM 008982.1, NM 008982.2, NM 008982.3,

NM 008982.4, NM 008982.5, BC148595, BC156870

UniProt ID: Q64455

Summary: Tyrosine phosphatase which dephosphorylates or contributes to the dephosphorylation of

CTNND1, FLT3, PDGFRB, MET, RET, KDR, LYN, SRC, MAPK1, MAPK3, EGFR, TJP1, OCLN, PIK3R1 and PIK3R2. Plays a role in cell adhesion, migration, proliferation and differentiation. Involved in vascular development. May be involved in the mechanism of contact inhibition of cell growth. Regulator of macrophage adhesion and spreading. Positively affects cell-matrix adhesion. Positive regulator of platelet activation and thrombosis. Negative regulator of cell

proliferation. Negative regulator of PDGF-stimulated cell migration; through

dephosphorylation of PDGFR. Positive regulator of endothelial cell survival, as well as of VEGF-induced SRC and AKT activation; through KDR dephosphorylation. Negative regulator of EGFR signaling pathway; through EGFR dephosphorylation. Enhances the barrier function of epithelial junctions during reassembly. Negatively regulates T-cell receptor (TCR) signaling. Upon T-cell TCR activation, it is up-regulated and excluded from the immunological synapses, while upon T-cell-antigen presenting cells (APC) disengagement, it is no longer excluded and

can dephosphorylate PLCG1 and LAT to down-regulate prolongation of signaling.

[UniProtKB/Swiss-Prot Function]





shRNA Design:

These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact techsupport@origene.com. If you need a special design or shRNA sequence, please utilize our custom shRNA service.

Performance Guaranteed: OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).