

Product datasheet for TL504421V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Copb1 Mouse shRNA Lentiviral Particle (Locus ID 70349)

Product data:

Product Type: shRNA Lentiviral Particles

Product Name: Copb1 Mouse shRNA Lentiviral Particle (Locus ID 70349)

Locus ID: 70349

Synonyms: 2610019B04Rik

Vector: pGFP-C-shLenti (TR30023)

Format: Lentiviral particles

Components: Copb1 - Mouse shRNA lentiviral particles (4 unique 29mer target-specific shRNA, 1 scramble

control), 0.5 ml each, >10^7 TU/ml.

RefSeq: <u>BC030837</u>, <u>NM 033370</u>, <u>NM 033370.1</u>, <u>NM 033370.2</u>, <u>NM 033370.3</u>

UniProt ID: Q9JIF7

Summary: The coatomer is a cytosolic protein complex that binds to dilysine motifs and reversibly

associates with Golgi non-clathrin-coated vesicles, which further mediate biosynthetic protein transport from the ER, via the Golgi up to the trans Golgi network. Coatomer complex is required for budding from Golgi membranes, and is essential for the retrograde Golgi-to-ER transport of dilysine-tagged proteins. In mammals, the coatomer can only be recruited by membranes associated to ADP-ribosylation factors (ARFs), which are small GTP-binding proteins; the complex also influences the Golgi structural integrity, as well as the processing, activity, and endocytic recycling of LDL receptors. Involved in the Golgi disassembly and reassembly processes during cell cycle. Involved in autophagy by playing a role in early endosome function. Plays a role in organellar compartmentalization of secretory compartments including endoplasmic reticulum (ER)-Golgi intermediate compartment (ERGIC), Golgi, trans-Golgi network (TGN) and recycling endosomes, and in biosynthetic transport of CAV1 (By similarity). Plays a functional role in facilitating the transport of kappatype opioid receptor mRNAs into axons and enhances translation of these proteins in cortical neurons. Required for limiting lipid storage in lipid droplets. Involved in lipid homeostasis by regulating the presence of perilipin family members PLIN2 and PLIN3 at the lipid droplet

surface and promoting the association of adipocyte triglyceride lipase (PNPLA2) with the lipid

droplet surface to mediate lipolysis.[UniProtKB/Swiss-Prot Function]





Copb1 Mouse shRNA Lentiviral Particle (Locus ID 70349) - TL504421V

shRNA Design:

Performance Guaranteed: These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact <u>techsupport@origene.com</u>. If you need a special design or shRNA sequence, please utilize our <u>custom shRNA service</u>.

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).