

Product datasheet for TL506650V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Pak2 Mouse shRNA Lentiviral Particle (Locus ID 224105)

Product data:

Product Type: shRNA Lentiviral Particles

Product Name: Pak2 Mouse shRNA Lentiviral Particle (Locus ID 224105)

Locus ID: 224105

Synonyms: 5330420P17Rik; A130002K10Rik; Al836325; D16Ertd269e; mKIAA4182; PAK-2

Vector: pGFP-C-shLenti (TR30023)

Format: Lentiviral particles

Components: Pak2 - Mouse shRNA lentiviral particles (4 unique 29mer target-specific shRNA, 1 scramble

control), 0.5 ml each, >10^7 TU/ml.

RefSeq: <u>BC086650, NM 177326, NM 177326.1, NM 177326.2, NM 177326.3, BC029195, BC043712,</u>

BC063068

UniProt ID: Q8CIN4

Summary: Serine/threonine protein kinase that plays a role in a variety of different signaling pathways

including cytoskeleton regulation, cell motility, cell cycle progression, apoptosis or

proliferation. Acts as downstream effector of the small GTPases CDC42 and RAC1. Activation

by the binding of active CDC42 and RAC1 results in a conformational change and a

subsequent autophosphorylation on several serine and/or threonine residues. Full-length PAK2 stimulates cell survival and cell growth. Phosphorylates MAPK4 and MAPK6 and activates the downstream target MAPKAPK5, a regulator of F-actin polymerization and cell migration. Phosphorylates JUN and plays an important role in EGF-induced cell proliferation.

Phosphorylates many other substrates including histone H4 to promote assembly of H3.3 and H4 into nucleosomes, BAD, ribosomal protein S6, or MBP. Additionally, associates with

ARHGEF7 and GIT1 to perform kinase-independent functions such as spindle orientation control during mitosis. On the other hand, apoptotic stimuli such as DNA damage lead to caspase-mediated cleavage of PAK2, generating PAK-2p34, an active p34 fragment that translocates to the nucleus and promotes cellular apoptosis involving the JNK signaling

pathway. Caspase-activated PAK2 phosphorylates MKNK1 and reduces cellular translation (By

similarity).[UniProtKB/Swiss-Prot Function]

shRNA Design: These shRNA constructs were designed against multiple splice variants at this gene locus. To

be certain that your variant of interest is targeted, please contact <u>techsupport@origene.com</u>. If you need a special design or shRNA sequence, please utilize our custom shRNA service.






Performance Guaranteed:

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.


For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).

Product images:



GFP signal was observed under microscope at 48 hours after transduction of TL506650B virus into HEK293 cells. TL506650B virus was prepared using lenti-shRNA TL506650B and [TR30037] packaging kit.





GFP signal was observed under microscope at 48 hours after transduction of [TL506650C] virus into HEK293 cells. [TL506650C] virus was prepared using lenti-shRNA [TL506650C] and [TR30037] packaging kit.