

## Product datasheet for **TL506877**

### Zcchc11 Mouse shRNA Plasmid (Locus ID 230594)

#### Product data:

Product Type:	shRNA Plasmids
Product Name:	Zcchc11 Mouse shRNA Plasmid (Locus ID 230594)
Locus ID:	230594
Synonyms:	6030404K05Rik; 9230115F04Rik; mKIAA0191; PPAPD3
Vector:	pGFP-C-shLenti (TR30023)
E. coli Selection:	Chloramphenicol (34 ug/ml)
Mammalian Cell Selection:	Puromycin
Format:	Lentiviral plasmids
Components:	Zcchc11 - Mouse, 4 unique 29mer shRNA constructs in lentiviral GFP vector(Gene ID = 230594). 5µg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pGFP-C-shLenti Vector, TR30021, included for free.
RefSeq:	<a href="#">NM_175472</a> , <a href="#">NM_175472.1</a> , <a href="#">NM_175472.2</a> , <a href="#">NM_175472.3</a> , <a href="#">BC150791</a> , <a href="#">BC024681</a> , <a href="#">BC038132</a> , <a href="#">BC038931</a> , <a href="#">BC145507</a>
UniProt ID:	<a href="#">B2RX14</a>



[View online »](#)

**Summary:**

Uridyltransferase that mediates the terminal uridylation of mRNAs with short (less than 25 nucleotides) poly(A) tails, hence facilitating global mRNA decay (PubMed:28792939). Essential for both oocyte maturation and fertility. Through 3' terminal uridylation of mRNA, sculpt, with TUT7, the maternal transcriptome by eliminating transcripts during oocyte growth (PubMed:28792939). Involved in microRNA (miRNA)-induced gene silencing through uridylation of deadenylated miRNA targets. Also functions as an integral regulator of microRNA biogenesis using 3 different uridylation mechanisms (By similarity). Acts as a suppressor of miRNA biogenesis by mediating the terminal uridylation of some miRNA precursors, including that of let-7 (pre-let-7), miR107, miR-143 and miR-200c. Uridylated miRNAs are not processed by Dicer and undergo degradation. Degradation of pre-let-7 contributes to the maintenance of embryonic stem (ES) cell pluripotency (By similarity). Also catalyzes the 3' uridylation of miR-26A, a miRNA that targets IL6 transcript. This abrogates the silencing of IL6 transcript, hence promoting cytokine expression (PubMed:19703396). In the absence of LIN28A, TUT7 and TUT4 monouridylate group II pre-miRNAs, which includes most of pre-let7 members, that shapes an optimal 3' end overhang for efficient processing (PubMed:28671666). Add oligo-U tails to truncated pre-miRNAs with a 5' overhang which may promote rapid degradation of non-functional pre-miRNA species (By similarity). May also suppress Toll-like receptor-induced NF-kappa-B activation via binding to T2BP (By similarity). Does not play a role in replication-dependent histone mRNA degradation (By similarity). Due to functional redundancy between TUT4 and TUT7, the identification of the specific role of each of these proteins is difficult (PubMed:28671666, PubMed:28792939, PubMed:22898984). TUT4 and TUT7 restrict retrotransposition of long interspersed element-1 (LINE-1) in cooperation with MOV10 counteracting the RNA chaperone activity of L1RE1. TUT7 uridylates LINE-1 mRNAs in the cytoplasm which inhibits initiation of reverse transcription once in the nucleus, whereas uridylation by TUT4 destabilizes mRNAs in cytoplasmic ribonucleoprotein granules (By similarity).[UniProtKB/Swiss-Prot Function]

**shRNA Design:**

These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact [techsupport@origene.com](mailto:techsupport@origene.com). If you need a special design or shRNA sequence, please utilize our [custom shRNA service](#).

**Performance  
Guaranteed:**

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at [techsupport@origene.com](mailto:techsupport@origene.com). Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).