

Product datasheet for TL507515

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Atg9a Mouse shRNA Plasmid (Locus ID 245860)

Product data:

Product Type: shRNA Plasmids

Product Name: Atg9a Mouse shRNA Plasmid (Locus ID 245860)

Locus ID: 245860

Synonyms: Apg9l1; Atg9; Atg9l1; AU019532

Vector: pGFP-C-shLenti (TR30023)

E. coli Selection: Chloramphenicol (34 ug/ml)

Mammalian Cell

Selection:

Puromycin

Format: Lentiviral plasmids

Components: Atg9a - Mouse, 4 unique 29mer shRNA constructs in lentiviral GFP vector(Gene ID = 245860).

5µg purified plasmid DNA per construct

29-mer scrambled shRNA cassette in pGFP-C-shLenti Vector, TR30021, included for free.

RefSeq: BC079884, NM 001003917, NM 001288612, NM 001288613, NR 109938, NM 001003917.1,

NM 001003917.2, NM 001003917.3, NM 001003917.4, NM 001288613.1, NM 001288612.1,

BC034139, BC056488, BC082307

UniProt ID: Q68FE2

Summary: Involved in autophagy and cytoplasm to vacuole transport (Cvt) vesicle formation. Plays a key

role in the organization of the preautophagosomal structure/phagophore assembly site (PAS), the nucleating site for formation of the sequestering vesicle. Cycles between a juxta-nuclear

trans-Golgi network compartment and late endosomes. Nutrient starvation induces

accumulation on autophagosomes. Starvation-dependent trafficking requires ULK1, ATG13 and SUPT20H (By similarity). Required for carbonyl cyanide m-chlorophenylhydrazone (CCCP)-induced ATG8 family proteins lipidation, a key autophagy step.[UniProtKB/Swiss-Prot

Function1

shRNA Design: These shRNA constructs were designed against multiple splice variants at this gene locus. To

be certain that your variant of interest is targeted, please contact techsupport@origene.com. If you need a special design or shRNA sequence, please utilize our custom shRNA service.





Performance Guaranteed:

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).