

## Product datasheet for **TL508190V**

### **H3c11 Mouse shRNA Lentiviral Particle (Locus ID 319153)**

#### **Product data:**

|                      |   |
|----------------------|---|
| <b>Product Type:</b> | shRNA Lentiviral Particles  |
| <b>Locus ID:</b>     | 319153  |
| <b>Synonyms:</b>     | H3-H; H3c1; H3c8; H3c10; Hist1; Hist1h3i  |
| <b>Vector:</b>       | pGFP-C-shLenti (TR30023)  |
| <b>Format:</b>       | Lentiviral particles  |
| <b>Components:</b>   | Hist1h3i - Mouse shRNA lentiviral particles (4 unique 29mer target-specific shRNA, 1 scramble control), 0.5 ml each, >10 <sup>7</sup> TU/ml.  |
| <b>RefSeq:</b>       | <a href="#">BC116383</a> , <a href="#">NM_178207</a> , <a href="#">NM_178207.1</a> , <a href="#">NM_178207.2</a> , <a href="#">BC107287</a> , <a href="#">BC127148</a>  |
| <b>UniProt ID:</b>   | <a href="#">P68433</a>  |
| <b>Summary:</b>      | Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. [provided by RefSeq, Aug 2015] |
| <b>shRNA Design:</b> | These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact <a href="mailto:techsupport@origene.com">techsupport@origene.com</a> . If you need a special design or shRNA sequence, please utilize our <a href="#">custom shRNA service</a> .  |



**Performance Guaranteed:** OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at [techsupport@origene.com](mailto:techsupport@origene.com). Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).