

## **Product datasheet for TL509328**

## **Alk Mouse shRNA Plasmid (Locus ID 11682)**

**Product data:** 

**Product Type:** shRNA Plasmids

**Product Name:** Alk Mouse shRNA Plasmid (Locus ID 11682)

**Locus ID:** 11682

Synonyms: CD246; Tcrz

**Vector:** pGFP-C-shLenti (TR30023)

E. coli Selection: Chloramphenicol (34 ug/ml)

**Mammalian Cell** 

Selection:

Puromycin

Format: Lentiviral plasmids

Components: Alk - Mouse, 4 unique 29mer shRNA constructs in lentiviral GFP vector(Gene ID = 11682). 5µg

purified plasmid DNA per construct

29-mer scrambled shRNA cassette in pGFP-C-shLenti Vector, TR30021, included for free.

**RefSeq:** NM 007439, NM 007439.1, NM 007439.2, BC169280, BC169281, BC169282, BC169283

UniProt ID: P97793

**Summary:** Neuronal receptor tyrosine kinase that is essentially and transiently expressed in specific

regions of the central and peripheral nervous systems and plays an important role in the genesis and differentiation of the nervous system. Transduces signals from ligands at the cell surface, through specific activation of the mitogen-activated protein kinase (MAPK) pathway. Phosphorylates almost exclusively at the first tyrosine of the Y-x-x-x-Y-Y motif. Following activation by ligand, ALK induces tyrosine phosphorylation of CBL, FRS2, IRS1 and SHC1, as well as of the MAP kinases MAPK1/ERK2 and MAPK3/ERK1. Acts as a receptor for ligands pleiotrophin (PTN), a secreted growth factor, and midkine (MDK), a PTN-related factor, thus participating in PTN and MDK signal transduction. PTN-binding induces MAPK pathway activation, which is important for the anti-apoptotic signaling of PTN and regulation of cell proliferation. MDK-binding induces phosphorylation of the ALK target insulin receptor substrate (IRS1), activates mitogen-activated protein kinases (MAPKs) and PI3-kinase, resulting also in cell proliferation induction. Drives NF-kappa-B activation, probably through IRS1 and the activation of the AKT serine/threonine kinase. Recruitment of IRS1 to activated ALK and the activation of NF-kappa-B are essential for the autocrine growth and survival signaling of

MDK.[UniProtKB/Swiss-Prot Function]



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



shRNA Design:

These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact <a href="mailto:techsupport@origene.com">techsupport@origene.com</a>. If you need a special design or shRNA sequence, please utilize our <a href="mailto:custom shRNA service">custom shRNA service</a>.

Performance Guaranteed: OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).