

Product datasheet for TL509787V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Ikbkb Mouse shRNA Lentiviral Particle (Locus ID 16150)

Product data:

Product Type: shRNA Lentiviral Particles

Product Name: Ikbkb Mouse shRNA Lentiviral Particle (Locus ID 16150)

Locus ID: 16150

Synonyms: Al132552; IKK-2; IKK-beta; IKK2; IKKbeta; IKK[b]

Vector: pGFP-C-shLenti (TR30023)

Format: Lentiviral particles

Components: Ikbkb - Mouse shRNA lentiviral particles (4 unique 29mer target-specific shRNA, 1 scramble

control), 0.5 ml each, >10^7 TU/ml.

RefSeq: <u>BC037723</u>, <u>NM 001159774</u>, <u>NM 010546</u>, <u>NM 001159774.1</u>, <u>NM 010546.1</u>, <u>BC004772</u>,

BC019210, BC020334

UniProt ID: 088351

Summary: Serine kinase that plays an essential role in the NF-kappa-B signaling pathway which is

activated by multiple stimuli such as inflammatory cytokines, bacterial or viral products, DNA

damages or other cellular stresses. Acts as part of the canonical IKK complex in the

conventional pathway of NF-kappa-B activation and phosphorylates inhibitors of NF-kappa-B on 2 critical serine residues. These modifications allow polyubiquitination of the inhibitors and subsequent degradation by the proteasome. In turn, free NF-kappa-B is translocated into

the nucleus and activates the transcription of hundreds of genes involved in immune response, growth control, or protection against apoptosis. In addition to the NF-kappa-B inhibitors, phosphorylates several other components of the signaling pathway including NEMO/IKBKG, NF-kappa-B subunits RELA and NFKB1, as well as IKK-related kinases TBK1 and IKBKE. IKK-related kinase phosphorylations may prevent the overproduction of inflammatory mediators since they exert a negative regulation on canonical IKKs. Phosphorylates FOXO3, mediating the TNF-dependent inactivation of this pro-apoptotic transcription factor. Also phosphorylates other substrates including NCOA3, BCL10 and IRS1. Within the nucleus, acts

as an adapter protein for NFKBIA degradation in UV-induced NF-kappa-B activation.

[UniProtKB/Swiss-Prot Function]

shRNA Design: These shRNA constructs were designed against multiple splice variants at this gene locus. To

be certain that your variant of interest is targeted, please contact <u>techsupport@origene.com</u>. If you need a special design or shRNA sequence, please utilize our <u>custom shRNA service</u>.







Performance Guaranteed:

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).