

Product datasheet for TL510042V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Fads1 Mouse shRNA Lentiviral Particle (Locus ID 76267)

Product data:

Product Type: shRNA Lentiviral Particles

Product Name: Fads1 Mouse shRNA Lentiviral Particle (Locus ID 76267)

Locus ID: 76267

Synonyms: 0710001O03Rik; A930006B21Rik; Al317215; DSD

Vector: pGFP-C-shLenti (TR30023)

Format: Lentiviral particles

Components: Fads1 - Mouse shRNA lentiviral particles (4 unique 29mer target-specific shRNA, 1 scramble

control), 0.5 ml each, >10^7 TU/ml.

RefSeq: <u>BC022139</u>, <u>BC026831</u>, <u>BC026848</u>, <u>BC063053</u>, <u>NM 146094</u>, <u>NM 146094.1</u>, <u>NM 146094.2</u>,

BC013226

UniProt ID: Q920L1

Summary: Acts as a front-end fatty acyl-coenzyme A (CoA) desaturase that introduces a cis double bond

at carbon 5 located between a preexisting double bond and the carboxyl end of the fatty acyl chain. Involved in biosynthesis of highly unsaturated fatty acids (HUFA) from the essential polyunsaturated fatty acids (PUFA) linoleic acid (LA) (18:2n-6) and alpha-linolenic acid (ALA) (18:3n-3) precursors. Specifically, desaturates dihomo-gamma-linoleoate (DGLA) (20:3n-6) and

eicosatetraenoate (ETA) (20:4n-3) to generate arachidonate (AA) (20:4n-6) and

eicosapentaenoate (EPA) (20:5n-3), respectively (Probable). As a rate limiting enzyme for DGLA (20:3n-6) and AA (20:4n-6)-derived eicosanoid biosynthesis, controls the metabolism of

inflammatory lipids like prostaglandin E2, critical for efficient acute inflammatory response and maintenance of epithelium homeostasis. Contributes to membrane phospholipid biosynthesis by providing AA (20:4n-6) as a major acyl chain esterified into phospholipids. In particular, regulates phosphatidylinositol-4,5-bisphosphate levels, modulating inflammatory cytokine production in T-cells (PubMed:22534642). Also desaturates (11E)-octadecenoate (trans-vaccenoate)(18:1n-9), a metabolite in the biohydrogenation pathway of LA (18:2n-6) (By

similarity).[UniProtKB/Swiss-Prot Function]

shRNA Design: These shRNA constructs were designed against multiple splice variants at this gene locus. To

be certain that your variant of interest is targeted, please contact <u>techsupport@origene.com</u>. If you need a special design or shRNA sequence, please utilize our <u>custom shRNA service</u>.







Performance Guaranteed:

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).