

Product datasheet for **TL511715V**

Spast Mouse shRNA Lentiviral Particle (Locus ID 50850)

Product data:

Product Type:	shRNA Lentiviral Particles
Product Name:	Spast Mouse shRNA Lentiviral Particle (Locus ID 50850)
Locus ID:	50850
Synonyms:	mKIAA1083; Spg4
Vector:	pGFP-C-shLenti (TR30023)
Format:	Lentiviral particles
Components:	Spast - Mouse shRNA lentiviral particles (4 unique 29mer target-specific shRNA, 1 scramble control), 0.5 ml each, >10 ⁷ TU/ml.
RefSeq:	BC046286 , NM_001162870 , NM_016962 , NM_001357738 , NM_016962.1 , NM_016962.2 , NM_001162870.1
UniProt ID:	Q9QYY8
Summary:	ATP-dependent microtubule severing protein that specifically recognizes and cuts microtubules that are polyglutamylated (PubMed:19141076 PubMed:20530212). Preferentially recognizes and acts on microtubules decorated with short polyglutamate tails: severing activity increases as the number of glutamates per tubulin rises from one to eight, but decreases beyond this glutamylation threshold (By similarity). Severing activity is not dependent on tubulin acetylation or detyrosination (By similarity). Microtubule severing promotes reorganization of cellular microtubule arrays and the release of microtubules from the centrosome following nucleation (By similarity). It is critical for the biogenesis and maintenance of complex microtubule arrays in axons, spindles and cilia (By similarity). SPAST is involved in abscission step of cytokinesis and nuclear envelope reassembly during anaphase in cooperation with the ESCRT-III complex (By similarity). Recruited at the midbody, probably by IST1, and participates in membrane fission during abscission together with the ESCRT-III complex (By similarity). Recruited to the nuclear membrane by IST1 and mediates microtubule severing, promoting nuclear envelope sealing and mitotic spindle disassembly during late anaphase (By similarity). Required for membrane traffic from the endoplasmic reticulum (ER) to the Golgi and endosome recycling (By similarity). Recruited by IST1 to endosomes and regulates early endosomal tubulation and recycling by mediating microtubule severing (By similarity). Probably plays a role in axon growth and the formation of axonal branches (PubMed:18234839).[UniProtKB/Swiss-Prot Function]



[View online »](#)

shRNA Design:

These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact techsupport@origene.com. If you need a special design or shRNA sequence, please utilize our [custom shRNA service](#).

Performance Guaranteed:

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).