

Product datasheet for TL512166

Sdcbp Mouse shRNA Plasmid (Locus ID 53378)

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Locus ID:53378Synonyms:MDA-9; Sycl; syntenin-1Vector:pGFP-C-shLenti (TR300)E. coli Selection:Chloramphenicol (34 u)Mammalian CellPuromycinSelection:Format:Format:Lentiviral plasmidsComponents:SdCbp - Mouse, 4 uniqu Sµg purified plasmid D 29-mer scrambled shRiRefSeq:BC019400, NM 001098UniProt ID:O08992Summary:Multifunctional adapte transmembrane protei tumorigenesis. Positive epithelial-to-mesenchy increase TGFB1 signalir interaction between TG degradation of TGFR1. (By similarity). Regulate variety of cancer types syndecans to cytoskele factor SOX4 to the IL-5 be required for the targ similarity).[UniProtKB/S	
Locus ID:53378Synonyms:MDA-9; Sycl; syntenin-1Vector:pGFP-C-shLenti (TR300)E. coli Selection:Chloramphenicol (34 u)Mammalian CellPuromycinSelection:Format:Format:Lentiviral plasmidsComponents:SdCbp - Mouse, 4 uniqu Sµg purified plasmid D 29-mer scrambled shRitRefSeq:BC019400, NM 001098UniProt ID:O08992Summary:Multifunctional adapte transmembrane protei tumorigenesis. Positive epithelial-to-mesenchy increase TGFB1 signalir interaction between TG degradation of TGFR1. (By similarity). Regulate variety of cancer types syndecans to cytoskele factor SOX4 to the IL-5 be required for the targ similarity).[UniProtKB/S	
Synonyms:MDA-9; Sycl; syntenin-1Vector:pGFP-C-shLenti (TR300)E. coli Selection:Chloramphenicol (34 u)Mammalian CellPuromycinSelection:Format:Format:Lentiviral plasmidsComponents:Sdcbp - Mouse, 4 uniqueSyg purified plasmid D29-mer scrambled shR4RefSeq:BC019400, NM 001098UniProt ID:O08992Summary:Multifunctional adapte transmembrane proteit tumorigenesis. Positive epithelial-to-mesenchy increase TGFB1 signalir interaction between TG degradation of TGFR1. (By similarity). Regulate variety of cancer types syndecans to cytoskele factor SOX4 to the IL-5 be required for the targ similarity).[UniProtKB/S	asmid (Locus ID 53378)
Vector:pGFP-C-shLenti (TR300)E. coli Selection:Chloramphenicol (34 u)Mammalian CellPuromycinSelection:Format:Format:Lentiviral plasmidsComponents:Sdcbp - Mouse, 4 unique 5µg purified plasmid D 29-mer scrambled shRlRefSeq:BC019400, NM 001098UniProt ID:O08992Summary:Multifunctional adapte transmembrane protei tumorigenesis. Positive epithelial-to-mesenchy increase TGFB1 signalir interaction between TG degradation of TGFR1. (By similarity). Regulate variety of cancer types syndecans to cytoskele factor SOX4 to the IL-5 be required for the targ similarity).[UniProtKB/S	
E. coli Selection:Chloramphenicol (34 u)Mammalian CellPuromycinSelection:PuromycinFormat:Lentiviral plasmidsComponents:Sdcbp - Mouse, 4 uniqu 5µg purified plasmid D 29-mer scrambled shRIRefSeq:BC019400, NM 001098UniProt ID:O08992Summary:Multifunctional adapte transmembrane protei tumorigenesis. Positive epithelial-to-mesenchy increase TGFB1 signalir interaction between TG degradation of TGFR1. (By similarity). Regulate variety of cancer types syndecans to cytoskele factor SOX4 to the IL-5 be required for the targ similarity). [UniProtKB/S	
Mammalian Cell Selection:PuromycinFormat:Lentiviral plasmidsComponents:Sdcbp - Mouse, 4 uniqu Sµg purified plasmid D 29-mer scrambled shRIRefSeq:BC019400, NM 001098UniProt ID:O08992Summary:Multifunctional adapte transmembrane protei tumorigenesis. Positive epithelial-to-mesenchy increase TGFB1 signalir interaction between TG degradation of TGFR1. (By similarity). Regulate variety of cancer types syndecans to cytoskele factor SOX4 to the IL-5 be required for the targ similarity). [UniProtKB/S	23)
Selection:Format:Lentiviral plasmidsComponents:Sdcbp - Mouse, 4 unique 5µg purified plasmid D 29-mer scrambled shRieRefSeq:BC019400, NM 001098UniProt ID:O08992Summary:Multifunctional adapte transmembrane protei tumorigenesis. Positive epithelial-to-mesenchy increase TGFB1 signalir interaction between TG degradation of TGFR1. (By similarity). Regulate variety of cancer types syndecans to cytoskele factor SOX4 to the IL-5 be required for the targ similarity).[UniProtKB/S	ʒ/ml)
Components:Sdcbp - Mouse, 4 unique Sµg purified plasmid D 29-mer scrambled shRiRefSeq:BC019400, NM 001098UniProt ID:O08992Summary:Multifunctional adapted transmembrane proteid tumorigenesis. Positived epithelial-to-mesenchy increase TGFB1 signaling interaction between TGC degradation of TGFR1. (By similarity). Regulated variety of cancer types syndecans to cytoskeled factor SOX4 to the IL-5 be required for the targe similarity).[UniProtKB/5]	
5µg purified plasmid D 29-mer scrambled shRlRefSeq:BC019400, NM 001098UniProt ID:O08992Summary:Multifunctional adapte transmembrane protei tumorigenesis. Positive epithelial-to-mesenchy increase TGFB1 signalir interaction between TG degradation of TGFR1. (By similarity). Regulate variety of cancer types syndecans to cytoskele factor SOX4 to the IL-5 be required for the targ similarity).[UniProtKB/S	
UniProt ID:O08992Summary:Multifunctional adapte transmembrane protei tumorigenesis. Positive epithelial-to-mesenchy increase TGFB1 signalir interaction between TG degradation of TGFR1. (By similarity). Regulate variety of cancer types syndecans to cytoskele factor SOX4 to the IL-5 be required for the targ similarity).[UniProtKB/S	e 29mer shRNA constructs in lentiviral GFP vector(Gene ID = 53378). NA per construct NA cassette in pGFP-C-shLenti Vector, TR30021, included for free.
Summary: Multifunctional adapte transmembrane protei tumorigenesis. Positive epithelial-to-mesenchy increase TGFB1 signalir interaction between TG degradation of TGFR1. (By similarity). Regulate variety of cancer types syndecans to cytoskele factor SOX4 to the IL-5 be required for the targ similarity).[UniProtKB/5]	227, <u>NM 016807</u> , <u>NM 001098227.1</u> , <u>NM 016807.1</u> , <u>NM 016807.2</u>
transmembrane protei tumorigenesis. Positive epithelial-to-mesenchy increase TGFB1 signalir interaction between TG degradation of TGFR1. (By similarity). Regulate variety of cancer types syndecans to cytoskele factor SOX4 to the IL-5 be required for the targ similarity).[UniProtKB/5	
shRNA Design: These shRNA construct	r protein involved in diverse array of functions including trafficking of ns, neuro and immunomodulation, exosome biogenesis, and ly regulates TGFB1-mediated SMAD2/3 activation and TGFB1-induced mal transition (EMT) and cell migration in various cell types. May ng by enhancing cell-surface expression of TGFR1 by preventing the FR1 and CAV1 and subsequent CAV1-dependent internalization and In concert with SDC1/4 and PDCD6IP, regulates exosome biogenesis s migration, growth, proliferation, and cell cycle progression in a (PubMed:26539120). In adherens junctions may function to couple tal proteins or signaling components. Seems to couple transcription receptor (IL5RA). May also play a role in vesicular trafficking. Seems to geting of TGFA to the cell surface in the early secretory pathway (By wiss-Prot Function]
	s were designed against multiple splice variants at this gene locus. To riant of interest is targeted, please contact <u>techsupport@origene.com</u> . sign or shRNA sequence, please utilize our <u>custom shRNA service</u> .



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE Sdcbp Mouse shRNA Plasmid (Locus ID 53378) – TL512166

Performance Guaranteed: OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US