

## Product datasheet for **TL513087V**

### **Pak4 Mouse shRNA Lentiviral Particle (Locus ID 70584)**

#### **Product data:**

<b>Product Type:</b>	shRNA Lentiviral Particles
<b>Product Name:</b>	Pak4 Mouse shRNA Lentiviral Particle (Locus ID 70584)
<b>Locus ID:</b>	70584
<b>Synonyms:</b>	5730488L07Rik; AW555722; mKIAA1142
<b>Vector:</b>	pGFP-C-shLenti (TR30023)
<b>Format:</b>	Lentiviral particles
<b>Components:</b>	Pak4 - Mouse shRNA lentiviral particles (4 unique 29mer target-specific shRNA, 1 scramble control), 0.5 ml each, >10 <sup>7</sup> TU/ml.
<b>RefSeq:</b>	<a href="#">BC048238</a> , <a href="#">NM_027470</a> , <a href="#">NM_027470.1</a> , <a href="#">NM_027470.2</a> , <a href="#">NM_027470.3</a> , <a href="#">BC030389</a>
<b>UniProt ID:</b>	<a href="#">Q8BTW9</a>
<b>Summary:</b>	<p>Serine/threonine protein kinase that plays a role in a variety of different signaling pathways including cytoskeleton regulation, cell migration, growth, proliferation or cell survival. Activation by various effectors including growth factor receptors or active CDC42 and RAC1 results in a conformational change and a subsequent autophosphorylation on several serine and/or threonine residues. Phosphorylates and inactivates the protein phosphatase SSH1, leading to increased inhibitory phosphorylation of the actin binding/depolymerizing factor cofilin. Decreased cofilin activity may lead to stabilization of actin filaments. Phosphorylates LIMK1, a kinase that also inhibits the activity of cofilin. Phosphorylates integrin beta5/ITGB5 and thus regulates cell motility. Phosphorylates ARHGEF2 and activates the downstream target RHOA that plays a role in the regulation of assembly of focal adhesions and actin stress fibers. Stimulates cell survival by phosphorylating the BCL2 antagonist of cell death BAD. Alternatively, inhibits apoptosis by preventing caspase-8 binding to death domain receptors in a kinase independent manner. Plays a role in cell-cycle progression by controlling levels of the cell-cycle regulatory protein CDKN1A and by phosphorylating RAN. [UniProtKB/Swiss-Prot Function]</p>
<b>shRNA Design:</b>	These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact <a href="mailto:techsupport@origene.com">techsupport@origene.com</a> . If you need a special design or shRNA sequence, please utilize our <a href="#">custom shRNA service</a> .




[View online >](#)


**Performance  
Guaranteed:**

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.


For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at [techsupport@origene.com](mailto:techsupport@origene.com). Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).

**Product images:**


GFP signal was observed under microscope at 48 hours after transduction of TL513087A virus into HEK293 cells. TL513087A virus was prepared using lenti-shRNA TL513087A and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of TL513087B virus into HEK293 cells. TL513087B virus was prepared using lenti-shRNA TL513087B and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of [TL513087C] virus into HEK293 cells. [TL513087C] virus was prepared using lenti-shRNA [TL513087C] and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of [TL513087D] virus into HEK293 cells. [TL513087D] virus was prepared using lenti-shRNA [TL513087D] and [TR30037] packaging kit.