

## Product datasheet for **TL514763V**

### Ar Mouse shRNA Lentiviral Particle (Locus ID 11835)

#### Product data:

Product Type:	shRNA Lentiviral Particles
Product Name:	Ar Mouse shRNA Lentiviral Particle (Locus ID 11835)
Locus ID:	11835
Synonyms:	AW320017; Tfm
Vector:	pGFP-C-shLenti (TR30023)
Format:	Lentiviral particles
Components:	Ar - Mouse shRNA lentiviral particles (4 unique 29mer target-specific shRNA, 1 scramble control), 0.5 ml each, >10 <sup>7</sup> TU/ml.
RefSeq:	<a href="#">NM_013476</a> , <a href="#">NM_013476.1</a> , <a href="#">NM_013476.2</a> , <a href="#">NM_013476.3</a> , <a href="#">NM_013476.4</a> , <a href="#">BC148496</a> , <a href="#">BC153098</a>
UniProt ID:	<a href="#">P19091</a>
Summary:	This gene encodes a nuclear hormone receptor containing zinc finger and DNA-binding domains. The encoded protein is a key regulator of signalling by androgens, a class of steroid hormones involved in male reproductive development. The protein responds to hormone signalling by translocating to the nucleus, forming dimers, and binding to androgen response elements (AREs) in the promoters of target genes, which are subsequently transcriptionally activated. Activity of this protein is negatively regulated by nuclear receptor subfamily 0 group B member 1 (Nr0b1, also known as Dax1). Mutations in this gene result in feminized genitals and infertility in male animals. Loss of function in female animals also causes problems in reproductive development and function. [provided by RefSeq, May 2015]
shRNA Design:	These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact <a href="mailto:techsupport@origene.com">techsupport@origene.com</a> . If you need a special design or shRNA sequence, please utilize our <a href="#">custom shRNA service</a> .




[View online »](#)


**Performance  
Guaranteed:**

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.


For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at [techsupport@origene.com](mailto:techsupport@origene.com). Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).

**Product images:**


GFP signal was observed under microscope at 48 hours after transduction of TL514763A virus into HEK293 cells. TL514763A virus was prepared using lenti-shRNA TL514763A and [TR30037] packaging kit.




GFP signal was observed under microscope at 48 hours after transduction of TL514763A virus into HEK293 cells. TL514763A virus was prepared using lenti-shRNA TL514763A and [TR30037] packaging kit.




GFP signal was observed under microscope at 48 hours after transduction of TL514763B virus into HEK293 cells. TL514763B virus was prepared using lenti-shRNA TL514763B and [TR30037] packaging kit.




GFP signal was observed under microscope at 48 hours after transduction of TL514763B virus into HEK293 cells. TL514763B virus was prepared using lenti-shRNA TL514763B and [TR30037] packaging kit.




GFP signal was observed under microscope at 48 hours after transduction of [TL514763C] virus into HEK293 cells. [TL514763C] virus was prepared using lenti-shRNA [TL514763C] and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of [TL514763C] virus into HEK293 cells. [TL514763C] virus was prepared using lenti-shRNA [TL514763C] and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of [TL514763D] virus into HEK293 cells. [TL514763D] virus was prepared using lenti-shRNA [TL514763D] and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of [TL514763D] virus into HEK293 cells. [TL514763D] virus was prepared using lenti-shRNA [TL514763D] and [TR30037] packaging kit.