

## Product datasheet for **TL517589**

### Card9 Mouse shRNA Plasmid (Locus ID 332579)

#### Product data:

Product Type:	shRNA Plasmids
Product Name:	Card9 Mouse shRNA Plasmid (Locus ID 332579)
Locus ID:	332579
Synonyms:	Gm782
Vector:	pGFP-C-shLenti (TR30023)
E. coli Selection:	Chloramphenicol (34 ug/ml)
Mammalian Cell Selection:	Puromycin
Format:	Lentiviral plasmids
Components:	Card9 - Mouse, 4 unique 29mer shRNA constructs in lentiviral GFP vector(Gene ID = 332579). 5µg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pGFP-C-shLenti Vector, TR30021, included for free.
RefSeq:	<a href="#">NM_001037747</a> , <a href="#">NM_001037747.1</a> , <a href="#">BC065797</a> , <a href="#">BC151023</a> , <a href="#">BC166001</a> , <a href="#">NM_001037747.3</a>
UniProt ID:	<a href="#">A2AIV8</a>



[View online »](#)

<b>Summary:</b>	<p>Adapter protein that plays a key role in innate immune response to a number of intracellular pathogens, such as <i>C.albicans</i> and <i>L.monocytogenes</i>. Is at the crossroads of ITAM-tyrosine kinase and the Toll-like receptors (TLR) and NOD2 signaling pathways (PubMed:17514206). Probably controls various innate immune response pathways depending on the intracellular pathogen. Controls CLEC7A (dectin-1)-mediated myeloid cell activation induced by the yeast cell wall component zymosan, leading to cytokine production and innate anti-fungal immunity: acts by regulating BCL10-MALT1-mediated NF-kappa-B activation pathway. Activates NF-kappa-B via BCL10 (PubMed:16862125). In response to the hyphal form of <i>C.albicans</i>, mediates CLEC6A (dectin-2)-induced I-kappa-B kinase ubiquitination, leading to NF-kappa-B activation via interaction with BCL10 (PubMed:20538615). In response to <i>L.monocytogenes</i> infection, acts by connecting NOD2 recognition of peptidoglycan to downstream activation of MAP kinases (MAPK) without activating NF-kappa-B (PubMed:17187069). In response to fungal infection, may be required for the development and subsequent differentiation of interleukin 17-producing T helper (TH-17) cells (PubMed:17450144). Also involved in activation of myeloid cells via classical ITAM-associated receptors and TLR: required for TLR-mediated activation of MAPK, while it is not required for TLR-induced activation of NF-kappa-B (PubMed:17486093).[UniProtKB/Swiss-Prot Function]</p>
<b>shRNA Design:</b>	<p>These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact <a href="mailto:techsupport@origene.com">techsupport@origene.com</a>. If you need a special design or shRNA sequence, please utilize our <a href="#">custom shRNA service</a>.</p>
<b>Performance Guaranteed:</b>	<p>OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.</p> <p>For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at <a href="mailto:techsupport@origene.com">techsupport@origene.com</a>. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).</p>