

## Product datasheet for **TL519626V**

### Kdm1a Mouse shRNA Lentiviral Particle (Locus ID 99982)


#### Product data:

Product Type:	shRNA Lentiviral Particles
Product Name:	Kdm1a Mouse shRNA Lentiviral Particle (Locus ID 99982)
Locus ID:	99982
Synonyms:	1810043O07Rik; AA408884; Aof2; D4Ertd478e; Kdm1; Lsd1; mKIAA0601
Vector:	pGFP-C-shLenti (TR30023)
Format:	Lentiviral particles
Components:	Aof2 - Mouse shRNA lentiviral particles (4 unique 29mer target-specific shRNA, 1 scramble control), 0.5 ml each, >10 <sup>7</sup> TU/ml.
RefSeq:	<a href="#">BC059885</a> , <a href="#">NM_001347221</a> , <a href="#">NM_133872</a> , <a href="#">NM_001356567</a> , <a href="#">NM_133872.1</a> , <a href="#">NM_133872.2</a> , <a href="#">BC019417</a>
UniProt ID:	<a href="#">Q6ZQ88</a>
Summary:	<p>Histone demethylase that can demethylate both 'Lys-4' (H3K4me) and 'Lys-9' (H3K9me) of histone H3, thereby acting as a coactivator or a corepressor, depending on the context. Acts by oxidizing the substrate by FAD to generate the corresponding imine that is subsequently hydrolyzed. Acts as a corepressor by mediating demethylation of H3K4me, a specific tag for epigenetic transcriptional activation. Demethylates both mono- (H3K4me1) and di-methylated (H3K4me2) H3K4me. May play a role in the repression of neuronal genes. Alone, it is unable to demethylate H3K4me on nucleosomes and requires the presence of RCOR1/CoREST to achieve such activity. Also acts as a coactivator of androgen receptor (ANDR)-dependent transcription, by being recruited to ANDR target genes and mediating demethylation of H3K9me, a specific tag for epigenetic transcriptional repression. The presence of PRKCB in ANDR-containing complexes, which mediates phosphorylation of 'Thr-6' of histone H3 (H3T6ph), a specific tag that prevents demethylation H3K4me, prevents H3K4me demethylase activity of KDM1A. Demethylates di-methylated 'Lys-370' of p53/TP53 which prevents interaction of p53/TP53 with TP53BP1 and represses p53/TP53-mediated transcriptional activation (By similarity). Demethylates and stabilizes the DNA methylase DNMT1. Required for gastrulation during embryogenesis. Component of a RCOR/GFI/KDM1A/HDAC complex that suppresses, via histone deacetylase (HDAC) recruitment, a number of genes implicated in multilineage blood cell development. Effector of SNAI1-mediated transcription repression of E-cadherin/CDH1, CDN7 and KRT8. Required for the maintenance of the silenced state of the SNAI1 target genes E-cadherin/CDH1 and CDN7.[UniProtKB/Swiss-Prot Function]</p>




[View online »](#)


- shRNA Design:** These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact [techsupport@origene.com](mailto:techsupport@origene.com). If you need a special design or shRNA sequence, please utilize our [custom shRNA service](#).
- Performance Guaranteed:** OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.
- For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at [techsupport@origene.com](mailto:techsupport@origene.com). Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).

**Product images:**


GFP signal was observed under microscope at 48 hours after transduction of TL519626A virus into HEK293 cells. TL519626A virus was prepared using lenti-shRNA TL519626A and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of TL519626B virus into HEK293 cells. TL519626B virus was prepared using lenti-shRNA TL519626B and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of [TL519626C] virus into HEK293 cells. [TL519626C] virus was prepared using lenti-shRNA [TL519626C] and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of [TL519626D] virus into HEK293 cells. [TL519626D] virus was prepared using lenti-shRNA [TL519626D] and [TR30037] packaging kit.