

## Product datasheet for **TP300053L**

### **POLDIP2 (NM\_015584) Human Recombinant Protein**

#### **Product data:**

<b>Product Type:</b>	Recombinant Proteins
<b>Description:</b>	Recombinant protein of human polymerase (DNA-directed), delta interacting protein 2 (POLDIP2), 1 mg
<b>Species:</b>	Human
<b>Expression Host:</b>	HEK293T
<b>Expression cDNA Clone or AA Sequence:</b>	>RC200053 protein sequence <b>Red</b> =Cloning site <b>Green</b> =Tags(s)

MAACTARRALAVGSRWWSRSLTGARWPRPLCAAAGAGAFSPASTTTTTRRHLSRNRPEGKVLETGVFEV  
PKQNGKYETGQLFLHSIFGYRGVLFQWQARLYDRDVAASAEKAENPAGHGSKEVKGKHTHTYYQVLIDA  
RDCPHISQRSQTEAVTFLANHDDSRALYAIPGLDYVSHEDILPYTSTDQVPIQHELFFLLYDQTKAPP  
FVARETLRAWQEKNHPWLELSDVHRETTENIRVTVIPFYMGMRQAQNSHVYWWRYCIRLENLDSDDVVQLR  
ERHWRIFSLSGTLETVRGRGVGREGPVLSKEQPAFYSSHVSLQASSGHMWGTFRFRPDGSHFDVRIIP  
FSLESNKDEKTPPSGLHW

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV**


<b>Tag:</b>	C-Myc/DDK
<b>Predicted MW:</b>	41.9 kDa
<b>Concentration:</b>	>0.05 µg/µL as determined by microplate BCA method
<b>Purity:</b>	> 80% as determined by SDS-PAGE and Coomassie blue staining
<b>Buffer:</b>	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
<b>Preparation:</b>	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
<b>Note:</b>	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
<b>Storage:</b>	Store at -80°C.
<b>Stability:</b>	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
<b>RefSeq:</b>	<u><a href="#">NP_056399</a></u>



[View online »](#)

Locus ID:	26073
UniProt ID:	<a href="#">Q9Y2S7</a>
RefSeq Size:	2753
Cytogenetics:	17q11.2
RefSeq ORF:	1104
Synonyms:	p38; PDIP38; POLD4
Summary:	This gene encodes a protein that interacts with the DNA polymerase delta p50 subunit, as well as with proliferating cell nuclear antigen. The encoded protein maybe play a role in the ability of the replication fork to bypass DNA lesions. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2014]

### Product images:



Coomassie blue staining of purified POLDIP2 protein (Cat# [TP300053]). The protein was produced from HEK293T cells transfected with POLDIP2 cDNA clone (Cat# [RC200053]) using MegaTran 2.0 (Cat# [TT210002]).