

Product datasheet for TP300085L

CINP (NM_032630) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins Description: Recombinant protein of human cyclin-dependent kinase 2-interacting protein (CINP), 1 mg Species: Human HEK293T **Expression Host:** Expression cDNA Clone >RC200085 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s) MEAKTLGTVTPRKPVLSVSARKIKDNAADWHNLILKWETLNDAGFTTANNIANLKISLLNKDKIELDSSS PASKENEEKVCLEYNEELEKLCEELQATLDGLTKIQVKMEKLSSTTKGICELENYHYGEESKRPPLFHTW PTTHFYEVSHKLLEMYRKELLLKRTVAKELAHTGDPDLTLSYLSMWLHQPYVESDSRLHLESMLLETGHR AL **TRTRPLEQKLISEEDLAANDILDYKDDDDKV** Tag: C-Myc/DDK Predicted MW: 24.1 kDa **Concentration:** >0.05 µg/µL as determined by microplate BCA method > 80% as determined by SDS-PAGE and Coomassie blue staining Purity: **Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol Recombinant protein was captured through anti-DDK affinity column followed by **Preparation:** conventional chromatography steps. Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process. Store at -80°C. Storage: Stable for 12 months from the date of receipt of the product under proper storage and Stability: handling conditions. Avoid repeated freeze-thaw cycles. **RefSeq:** NP 116019 51550 Locus ID: **UniProt ID:** Q9BW66, A0A024R6M9



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

	CINP (NM_032630) Human Recombinant Protein – TP300085L	
RefSeq Size:	996	
Cytogenetics:	14q32.31	
RefSeq ORF:	636	
Summary:	The protein encoded by this gene is reported to be a component of the DNA replication complex as well as a genome-maintenance protein. It may interact with proteins important for replication initiation and has been shown to bind chromatin at the G1 phase of the cell cycle and dissociate from chromatin with replication initiation. It may also serve to regulate checkpoint signaling as part of the DNA damage response. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2016]	
Protein Families	: Druggable Genome	

Product images:

188	_	
98	-	
62	_	
49	_	
38	_	_
28	_	
17 14 63	Ξ	

Coomassie blue staining of purified CINP protein (Cat# [TP300085]). The protein was produced from HEK293T cells transfected with CINP cDNA clone (Cat# [RC200085]) using MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US