

Product datasheet for **TP300131M**

NAD Synthetase (NADSYN1) (NM_018161) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human NAD synthetase 1 (NADSYN1), 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC200131 protein sequence Red=Cloning site Green=Tags(s)

MGRKVTVATCALNQWALDFEGLNLRILKSIEIAKNRGARYRLGPELEICGYGCWDHYYESDTLLHSFQVL
AALLESPTQDIICDVGMPVMHRNVRYNCRVIFLNRKILLIRPKMALANEGNYRELWFTPWRSRSTHTTE
YFLPRMIQDLTKQETVFPFGDAVLVTWDTICIGSEICEELWTPHSPHIDMGLDGEIITNASGSHHVLKANK
TRVDLVTMVTSKNGGIYLLANQKGCDDGRLYYDGCAMIAMNGSVFAQGSQFSLDDVEVLTATLDLEDVRS
YRAEISSRNLAASRASPYPYPRVKVDFALSCHEDLLAPISEPIEWKYHSPEEEISLGPACWLWDFLRRSQQA
GFLLPLSGGVDSAATACLIYSMCCQVCEAVRSGNEEVLADVRTIVNQISYTPQDPRDLCGRILTTCYMAS
KNSSQETCTRARELAQQIGSHHISLNIDPAVKAVMGIFSLVTGKSPLFAAHGSSRENALQNVQARIRM
VLAYLFAQLSLWSRGVHGGLLVLGSANVDESLLGYLTKYDCSSADINPIGGISKTDLRAFVQFCIQRFQL
PALQSILLAPATAELEPLADGQVSQTDEEDMGMTYAELSVYGKLRKVAKMGPYSMFCKLLGMWRHICTPR
QVADKVKRFFSKYSMNRHKMTTLTPAYHAENYSPEDNRFDLRPFYNTSWPWQFRCIENQVLQLERAEPQ
SLDGVD

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	79.1 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: [NP_060631](#)

Locus ID: 55191

UniProt ID: [Q6IA69](#)

RefSeq Size: 2453

Cytogenetics: 11q13.4


RefSeq ORF: 2118

Synonyms: VCRL3

Summary: Nicotinamide adenine dinucleotide (NAD) is a coenzyme in metabolic redox reactions, a precursor for several cell signaling molecules, and a substrate for protein posttranslational modifications. NAD synthetase (EC 6.3.5.1) catalyzes the final step in the biosynthesis of NAD from nicotinic acid adenine dinucleotide (NaAD).[supplied by OMIM, Apr 2004]

Protein Pathways: Metabolic pathways, Nicotinate and nicotinamide metabolism

Product images:



Coomassie blue staining of purified NADSYN1 protein (Cat# [TP300131]). The protein was produced from HEK293T cells transfected with NADSYN1 cDNA clone (Cat# [RC200131]) using MegaTran 2.0 (Cat# [TT210002]).