

Product datasheet for TP300240M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

UBE2C (NM_181801) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human ubiquitin-conjugating enzyme E2C (UBE2C), transcript variant 4,

100 µg

Species: Human Expression Host: HEK293T

Expression cDNA >RC200240 protein sequence Clone or AA Sequence: Red=Cloning site Green=Tags(s)

MASQNRDPAATSVAAARKGAEPSGGAARGPVGKRLQQELMTLMMSGDKGISAFPESDNLFKWVGTIHGAA

GTVYEDLRYKLSLEFPSGYPYNAPTVKFLTPCYHPNVDTQGNICLDILKEKWSALYDVRTILLSIQSLLG

EPNIDSPLNTHAAELWKNPTAFKKYLQETYSKQVTSQEP

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 15.6 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling

conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 861517

Locus ID: 11065

UniProt ID: 000762





RefSeq Size: 901

Cytogenetics: 20q13.12

RefSeq ORF: 540

Synonyms: dJ447F3.2; UBCH10

Summary: The modification of proteins with ubiquitin is an important cellular mechanism for targeting


abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, ubiquitin-conjugating enzymes, and ubiquitin-protein ligases. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. The encoded protein is required for the destruction of mitotic cyclins and for cell cycle progression, and may be involved in cancer progression. Multiple transcript variants encoding different isoforms have been found for this gene. Pseudogenes of this gene have been defined on

chromosomes 4, 14, 15, 18, and 19. [provided by RefSeq, Aug 2013]

Protein Families: Druggable Genome

Protein Pathways: Ubiquitin mediated proteolysis

Product images:



Coomassie blue staining of purified UBE2C protein (Cat# [TP300240]). The protein was produced from HEK293T cells transfected with UBE2C cDNA clone (Cat# [RC200240]) using MegaTran 2.0 (Cat# [TT210002]).