

Product datasheet for TP300251

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MRG15 (MORF4L1) (NM_006791) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human mortality factor 4 like 1 (MORF4L1), transcript variant 1, 20 μg

Species: Human Expression Host: HEK293T

Expression cDNA Clone or AA Sequence: >RC200251 protein sequence Red=Cloning site Green=Tags(s)

MAPKQDPKPKFQEGERVLCFHGPLLYEAKCVKVAIKDKQVKYFIHYSGWNKNWDEWVPESRVLKYVDTNL QKQRELQKANQEQYAEGKMRGAAPGKKTSGLQQKNVEVKTKKNKQKTPGNGDGGSTSETPQPPRKKRARV DPTVENEETFMNRVEVKVKIPEELKPWLVDDWDLITRQKQLFYLPAKKNVDSILEDYANYKKSRGNTDNK EYAVNEVVAGIKEYFNVMLGTQLLYKFERPQYAEILADHPDAPMSQVYGAPHLLRLFVRIGAMLAYTPLD

EKSLALLLNYLHDFLKYLAKNSATLFSASDYEVAPPEYHRKAV

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 37.1 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling

conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 006782

Locus ID: 10933



MRG15 (MORF4L1) (NM_006791) Human Recombinant Protein - TP300251

UniProt ID:Q9UBU8RefSeq Size:1894Cytogenetics:15q25.1RefSeq ORF:969

Synonyms: Eaf3; FWP006; HsT17725; MEAF3; MORFRG15; MRG15; S863-6

Summary: Component of the NuA4 histone acetyltransferase (HAT) complex which is involved in


transcriptional activation of select genes principally by acetylation of nucleosomal histones H4 and H2A. This modification may both alter nucleosome - DNA interactions and promote interaction of the modified histones with other proteins which positively regulate transcription. This complex may be required for the activation of transcriptional programs associated with oncogene and proto-oncogene mediated growth induction, tumor suppressor mediated growth arrest and replicative senescence, apoptosis, and DNA repair. The NuA4 complex ATPase and helicase activities seem to be, at least in part, contributed by the association of RUVBL1 and RUVBL2 with EP400. NuA4 may also play a direct role in DNA repair when directly recruited to sites of DNA damage. Also component of the mSin3A complex which acts to repress

transcription by deacetylation of nucleosomal histones. Required for homologous recombination repair (HRR) and resistance to mitomycin C (MMC). Involved in the localization of PALB2, BRCA2

and RAD51, but not BRCA1, to DNA-damage foci.[UniProtKB/Swiss-Prot Function]

Protein Families: Transcription Factors

Product images:



Coomassie blue staining of purified MORF4L1 protein (Cat# TP300251). The protein was produced from HEK293T cells transfected with MORF4L1 cDNA clone (Cat# [RC200251]) using MegaTran 2.0 (Cat# [TT210002]).