

## Product datasheet for **TP300279**

### RPC62 (POLR3C) (NM\_006468) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human polymerase (RNA) III (DNA directed) polypeptide C (62kD) (POLR3C), 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC200279 protein sequence <b>Red</b> =Cloning site <b>Green</b> =Tags(s)

MTQAEIKLCSLLLQEHFGEIVEKIGVHLIRTGSQPLRVIAHDTGTSLDQVKKALCVLVQHNLVSYQVHKR  
GVVEYEAQCSRVLRLRMLRYPRIYITTKLYSDTGELIVEELLNGKLTMSAVVKKVADRLTETMEDGKTMD  
YAEVSNFTVRLADTHFVQRCPSVPTTENS DP G P P P P A P T L V I N E K D M Y L V P K L S L I G K G R R R S S D E D A A  
GEPKAKRPKYTTDNKEPIPDGIIYQANLDRFHQHFRDQAI V S A V A N R M D Q T S S E I V R T M L R M S E I T T S S  
SAPFTQPLSSNEIFRSLPVGYNISKQVLDQYLTL L A D D P L E F V G K S G D S G G G M Y I N L H K A L A S L A T A T L  
ESVQERFGSRCARIFRLVLQKKHIEQKQVEDFAMIPAKEAKDMLYKMLSENFMSLQEIPKTPDHAPSRT  
FYLYTVNILSAARMLLHRCYKSANLIERRQFETKENRLL E K S Q R V E A I I A S M Q A T G A E E A Q L Q E I E E M  
ITAPERQQL E T L K R N V N K L D A S E I Q V D E T I F L L E S Y I E C T M K R Q

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV**


Tag:	C-Myc/DDK
Predicted MW:	60.4 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

<b>Stability:</b>	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
<b>RefSeq:</b>	<a href="#">NP_006459</a>
<b>Locus ID:</b>	10623
<b>UniProt ID:</b>	<a href="#">Q9BU14</a>
<b>RefSeq Size:</b>	1888
<b>Cytogenetics:</b>	1q21.1
<b>RefSeq ORF:</b>	1602
<b>Synonyms:</b>	C82; RPC3; RPC62
<b>Summary:</b>	DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates. Specific core component of RNA polymerase III which synthesizes small RNAs, such as 5S rRNA and tRNAs. May direct with other members of the subcomplex RNA Pol III binding to the TFIIB-DNA complex via the interactions between TFIIB and POLR3F. May be involved either in the recruitment and stabilization of the subcomplex within RNA polymerase III, or in stimulating catalytic functions of other subunits during initiation. Plays a key role in sensing and limiting infection by intracellular bacteria and DNA viruses. Acts as nuclear and cytosolic DNA sensor involved in innate immune response. Can sense non-self dsDNA that serves as template for transcription into dsRNA. The non-self RNA polymerase III transcripts, such as Epstein-Barr virus-encoded RNAs (EBERs) induce type I interferon and NF- Kappa-B through the RIG-I pathway. Preferentially binds single-stranded DNA (ssDNA) in a sequence-independent manner (PubMed:21358628).[UniProtKB/Swiss-Prot Function]
<b>Protein Families:</b>	Transcription Factors
<b>Protein Pathways:</b>	Cytosolic DNA-sensing pathway, Metabolic pathways, Purine metabolism, Pyrimidine metabolism, RNA polymerase

### Product images:



Coomassie blue staining of purified POLR3C protein (Cat# TP300279). The protein was produced from HEK293T cells transfected with POLR3C cDNA clone (Cat# [RC200279]) using MegaTran 2.0 (Cat# [TT210002]).