

Product datasheet for **TP300410**

CD70 (NM_001252) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human CD70 molecule (CD70), 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC200410 protein sequence Red =Cloning site Green =Tags(s)
	 MPEEGSGCSVRRRPYGCVLRAALVPLVAGLVICLVCIQRFQAQQLPLESLGWDVAELQLNHTGPQQ D PRLYWQGGPALGRSFLHGPELDKGQLRIHRDGIYMVHIQVTLAICSSTASRHHPTTLAVGICSPASRSI SLLRLSFHQGCTIASQRLTPLARGDTLCTNLTGTLPSRNTDETFQVQWVRP TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Predicted MW:	20.9 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Bioactivity:	In vitro binding assay (PMID: 25730144)
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_001243
Locus ID:	970



[View online »](#)

UniProt ID: [P32970](#)
RefSeq Size: 913
Cytogenetics: 19p13.3
RefSeq ORF: 579
Synonyms: CD27-L; CD27L; CD27LG; LPFS3; TNFSF7; TNLG8A

Summary: The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF27/CD27. It is a surface antigen on activated, but not on resting, T and B lymphocytes. It induces proliferation of costimulated T cells, enhances the generation of cytolytic T cells, and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation, cytotoxic function of natural killer cells, and immunoglobulin synthesis. [provided by RefSeq, Jul 2008]

Protein Families: ES Cell Differentiation/IPS, Transmembrane

Protein Pathways: Cytokine-cytokine receptor interaction

Product images:

Antibody	IgG			DuetMab		
	K_{on} ($M^{-1} s^{-1}$)	K_{off} (s^{-1})	K_D (nM)	K_{on} ($M^{-1} s^{-1}$)	K_{off} (s^{-1})	K_D (nM)
Parental (against CD70) ^a	2.2×10^5	5.1×10^{-3}	23	2.0×10^5	4.9×10^{-3}	25
Parental (against CD4) ^b	2.1×10^5	1.8×10^{-4}	0.8	2.8×10^5	2.6×10^{-4}	0.9
V _K Y94A ^b	2.2×10^5	3.1×10^{-4}	1.4	ND	ND	
V _K S93A ^b	2.2×10^5	3.6×10^{-4}	1.6	ND	ND	
V _H D97A ^b	1.6×10^5	6.4×10^{-4}	4.1	ND	ND	
V _K Y92A ^b	1.2×10^5	1.8×10^{-3}	15	2.0×10^5	1.9×10^{-3}	10
V _K Y91A ^b	1.7×10^5	4.2×10^{-3}	25	2.8×10^5	4.7×10^{-3}	17
V _K R95A ^b	3.0×10^5	1.6×10^{-2}	55	5.4×10^5	2.3×10^{-2}	42
V _K R95A+V _H D97A ^b	2.7×10^5	1.8×10^{-2}	65	5.6×10^5	3.5×10^{-2}	63
V _H Y99A ^b	2.8×10^5	2.0×10^{-2}	72	ND	ND	
V _K S93A+V _H Y99A ^b	2.7×10^5	2.0×10^{-2}	74	ND	ND	
V _K Y94A+V _H Y99A ^b	2.7×10^5	2.1×10^{-2}	77	5.2×10^5	3.6×10^{-2}	70
V _K Y92A+V _H Y99A ^b	ND	ND	ND ^c	ND	ND	
V _K Y91A+V _H Y99A ^b	ND	ND	ND ^c	ND	ND	
V _K R95A+V _H Y99A ^b	ND	ND	ND ^c	ND	ND	

The binding affinity values of IgG and DuetMab with amino acid residue substitutions of the heavy chain and kappa light chain variable domains and their combinations of some of the substitutions to CD4 and CD70 (OriGene TP300410), as measured using an Octet384 instrument. Figure cited from MAbs, PMID: 25730144

Kinetic measurements to soluble monomeric forms of CD4 and CD70 were carried out using an Octet384 instrument. The dissociation constants, K_D , were calculated as the ratio of K_{off}/K_{on} from a non-linear fit of the data

^a Binding measured against CD70

^b Binding measured against CD4

^c No measurable binding could be seen

ND: not determined



Coomassie blue staining of purified CD70 protein (Cat# TP300410). The protein was produced from HEK293T cells transfected with CD70 cDNA clone (Cat# [RC200410]) using MegaTran 2.0 (Cat# [TT210002]).