

## Product datasheet for **TP300415L**

### Sorbitol Dehydrogenase (SORD) (NM\_003104) Human Recombinant Protein

#### Product data:

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human sorbitol dehydrogenase (SORD), 1 mg

**Species:** Human

**Expression Host:** HEK293T

**Expression cDNA Clone or AA Sequence:** >RC200415 protein sequence  
**Red**=Cloning site **Green**=Tags(s)

MAAAKPNLNSLVHGPDLRLNYPPEPGPNEVLLRMHSVGICGSDVHYWEYGRIGNFIVKKPMVLGH  
EASGTVEKVGSSVKHLKPGDRVAIEPGAPRENDEFCKMGRYNLSPSIFFCATPPDDGNLCRFYKHNA AFC  
YKLPDNTVFEEGALIEPLSVGIHACRRGGVTLGHKVLVCGAGPIGMVTLVAKAMGAAQVWVTDLSATRL  
SKAKEIGADLVLQISKESPQEIARKVEGQLGCKPEVTI ECTGAEASIQAGIYATRS GGTLVLVGLGSEMT  
TVPLLHAAIREVDIKGVFRYCNTPVVAISMLASKSVNVKPLVTHRFPLEKALEAFETFKKGLGLKIMLK  
DPDQNP

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV**

**Tag:** C-Myc/DDK

**Predicted MW:** 38.1 kDa

**Concentration:** >0.05 µg/µL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.

**Storage:** Store at -80°C.

**Stability:** Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** [NP\\_003095](#)

**Locus ID:** 6652



[View online »](#)

UniProt ID: [Q00796](#)

RefSeq Size: 2813

Cytogenetics: 15q21.1

RefSeq ORF: 1071


Synonyms: HEL-S-95n; RDH; SDH; SORD1; SORDD; XDH

**Summary:** Sorbitol dehydrogenase (SORD; EC 1.1.1.14) catalyzes the interconversion of polyols and their corresponding ketoses, and together with aldose reductase (ALDR1; MIM 103880), makes up the sorbitol pathway that is believed to play an important role in the development of diabetic complications (summarized by Carr and Markham, 1995 [PubMed 8535074]). The first reaction of the pathway (also called the polyol pathway) is the reduction of glucose to sorbitol by ALDR1 with NADPH as the cofactor. SORD then oxidizes the sorbitol to fructose using NAD(+) cofactor.[supplied by OMIM, Jul 2010]

**Protein Families:** Druggable Genome

**Protein Pathways:** Fructose and mannose metabolism, Metabolic pathways

### Product images:



Coomassie blue staining of purified SORD protein (Cat# [TP300415]). The protein was produced from HEK293T cells transfected with SORD cDNA clone (Cat# [RC200415]) using MegaTran 2.0 (Cat# [TT210002]).