

Product datasheet for TP300449L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Nck beta (NCK2) (NM_001004720) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human NCK adaptor protein 2 (NCK2), transcript variant 2, 1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC200449 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

42.7 kDa

MTEEVIVIAKWDYTAQQDQELDIKKNERLWLLDDSKTWWRVRNAANRTGYVPSNYVERKNSLKKGSLVKN LKDTLGLGKTRRKTSARDASPTPSTDAEYPANGSGADRIYDLNIPAFVKFAYVAEREDELSLVKGSRVTV MEKCSDGWWRGSYNGQIGWFPSNYVLEEVDEAAAESPSFLSLRKGASLSNGQGSRVLHVVQTLYPFSSVT EEELNFEKGETMEVIEKPENDPEWWKCKNARGQVGLVPKNYVVVLSDGPALHPAHAPQISYTGPSSSGRF AGREWYYGNVTRHQAECALNERGVEGDFLIRDSESSPSDFSVSLKASGKNKHFKVQLVDNVYCIGQRRFH

TMDELVEHYKKAPIFTSEHGEKLYLVRALQ

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW:

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeg: NP 001004720

Locus ID: 8440



Nck beta (NCK2) (NM_001004720) Human Recombinant Protein - TP300449L

UniProt ID: <u>043639</u>, <u>A0A0S2Z4M6</u>

RefSeq Size: 2417 Cytogenetics: 2q12.2 RefSeq ORF: 1140

Synonyms: GRB4; NCKbeta

Summary: This gene encodes a member of the NCK family of adaptor proteins. The protein contains

three SH3 domains and one SH2 domain. The protein has no known catalytic function but has been shown to bind and recruit various proteins involved in the regulation of receptor protein tyrosine kinases. It is through these regulatory activities that this protein is believed to be involved in cytoskeletal reorganization. Alternate transcriptional splice variants, encoding


different isoforms, have been characterized. [provided by RefSeq, Jul 2008]

Protein Families: Druggable Genome

Protein Pathways: Axon guidance, ErbB signaling pathway, Pathogenic Escherichia coli infection, T cell receptor

signaling pathway

Product images:



Coomassie blue staining of purified NCK2 protein (Cat# [TP300449]). The protein was produced from HEK293T cells transfected with NCK2 cDNA clone (Cat# [RC200449]) using MegaTran 2.0 (Cat# [TT210002]).