

Product datasheet for TP300450L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NAPG (NM_003826) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human N-ethylmaleimide-sensitive factor attachment protein,

gamma (NAPG), 1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone

or AA Sequence:

>RC200450 protein sequence
Red=Cloning site Green=Tags(s)

MAAQKINEGLEHLAKAEKYLKTGFLKWKPDYDSAASEYGKAAVAFKNAKQFEQAKDACLREAVAHENNR

Α

LFHAAKAYEQAGMMLKEMQKLPEAVQLIEKASMMYLENGTPDTAAMALERAGKLIENVDPEKAVQLYQ

QΤ

ANVFENDERLRQAVELLGKASRLLVRGRRFDEAALSIQKEKNIYKEIENYPTCYKKTIAQVLVHLHRNDY VAAERCVRESYSIPGFNGSEDCAALEQLLEGYDQQDQDQVSDVCNSPLFKYMDNDYAKLGLSLVVPGGGI

KKKSPATPQAKPDGVTATAADEEEDEYSGGLC

TRTRPLEQKLISEEDLAANDILDYKDDDDK**V**

Tag: C-Myc/DDK
Predicted MW: 34.6 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.





RefSeq: NP 003817

Locus ID: 8774

 UniProt ID:
 Q99747

 RefSeq Size:
 3729

Cytogenetics: 18p11.22

RefSeq ORF: 936

Synonyms: GAMMASNAP

Summary: This gene encodes soluble NSF attachment protein gamma. The soluble NSF attachment


proteins (SNAPs) enable N-ethyl-maleimide-sensitive fusion protein (NSF) to bind to target

membranes. NSF and SNAPs appear to be general components of the intracellular

membrane fusion apparatus, and their action at specific sites of fusion must be controlled by SNAP receptors particular to the membranes being fused. The product of this gene mediates platelet exocytosis and controls the membrane fusion events of this process.[provided by

RefSeq, Dec 2008]

Product images:



Coomassie blue staining of purified NAPG protein (Cat# [TP300450]). The protein was produced from HEK293T cells transfected with NAPG cDNA clone (Cat# [RC200450]) using MegaTran 2.0 (Cat# [TT210002]).