

Product datasheet for TP300460

Hydroxysteroid (17 beta) Dehydrogenase 4 (HSD17B4) (NM_000414) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human hydroxysteroid (17-beta) dehydrogenase 4 (HSD17B4), 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC200460 protein sequence Red=Cloning site Green=Tags(s)

MGSPFRFDGRVVLVTGAGAGLGRAYALAF AER GALVVNDLGGDFKGVGKGS LAADKVV EIRRRGGKAV
ANYDSVEEGEKVVK TALDAFGRIDVVVNNAGILRDRSFARISDEDEDWDIIHRVHLRGSFQVTRAAWEHMKK
QKYGRIIMTSSASGIYGNFGQANYSAAKLGLLGLANSLAIEGRKSNHCNTIAPNAGSRMTQTVMPEDLV
EALKPEYVAPLVLWLCHE SCEENGGLFEVGAGWIGKLRWERTLGAIVRQKNHPMTPEAVKANWKKICDFE
NASKPQSIQESTGSIIEVLSKIDSEGGV SANHTSRATSTATSGFAGAIGQKLPPFSYAYTELEAIMYALG
VGASIKDPKDLKFIYEGSSDF SCLPTFGVIIGQKSMGGGLAEIPGLSINF AKVLHGEQYLELYKPLPRA
GKLKCEAVVADVLDK GSGWIIIMDVYSYSEKELICHNQFSLFLVSGGFGGKRTSDKV KVAVAIPNRPD
AVLTDTTSLNQAALYRLSGDWNPLHIDPNFASLAGFDK PILHGLCTFGFSARRVLQQFADNDVSRFKAIK
ARFAKPVYPGQTLQTEMWKEGNRIHFQTKVQETGDIVISNAYVDLAPTS GTS AKTPSEGGK LQSTFV FEE
IGRRLKDIGPEVVKVNAVFEWHITKGGNIGAKWTIDLKSGSGKVYQGP AKGAADTTIILSDEDFMEVVL
GKLDPQKAFFSGRLKARGNIMLSQKLQMILKDYAKL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	79.5 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.




[View online »](#)

Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_000405
Locus ID:	3295
UniProt ID:	P51659 , A0A0S2Z4J1 , B2R659
RefSeq Size:	2710
Cytogenetics:	5q23.1
RefSeq ORF:	2208
Synonyms:	DBP; MFE-2; MFP-2; MPF-2; PRLTS1; SDR8C1
Summary:	The protein encoded by this gene is a bifunctional enzyme that is involved in the peroxisomal beta-oxidation pathway for fatty acids. It also acts as a catalyst for the formation of 3-ketoacyl-CoA intermediates from both straight-chain and 2-methyl-branched-chain fatty acids. Defects in this gene that affect the peroxisomal fatty acid beta-oxidation activity are a cause of D-bifunctional protein deficiency (DBPD). An apparent pseudogene of this gene is present on chromosome 8. Multiple alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, May 2014]
Protein Families:	Druggable Genome
Protein Pathways:	Metabolic pathways, Primary bile acid biosynthesis

Product images:



HSD17B4 enzymatic activity with 75µM β-estradiol as a substrate, measured by NADH production (indicated by luminescence).



Coomassie blue staining of purified HSD17B4 protein (Cat# TP300460). The protein was produced from HEK293T cells transfected with HSD17B4 cDNA clone (Cat# [RC200460]) using MegaTran 2.0 (Cat# [TT210002]).