

## **Product datasheet for TP300494M**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CDK2 (NM\_001798) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human cyclin-dependent kinase 2 (CDK2), transcript variant 1, 100 μg

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** >RC200494 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MENFQKVEKIGEGTYGVVYKARNKLTGEVVALKKIRLDTETEGVPSTAIREISLLKELNHPNIVKLLDVI HTENKLYLVFEFLHQDLKKFMDASALTGIPLPLIKSYLFQLLQGLAFCHSHRVLHRDLKPQNLLINTEGA IKLADFGLARAFGVPVRTYTHEVVTLWYRAPEILLGCKYYSTAVDIWSLGCIFAEMVTRRALFPGDSEID QLFRIFRTLGTPDEVVWPGVTSMPDYKPSFPKWARQDFSKVVPPLDEDGRSLLSQMLHYDPNKRISAKAA

LAHPFFQDVTKPVPHLRL

**TRTRPL**EQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

**Predicted MW:** 33.7 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** NP 001789

**Locus ID:** 1017



UniProt ID: <u>P24941</u>, <u>A0A024RB77</u>

RefSeq Size: 2301 Cytogenetics: 12q13.2 RefSeq ORF: 894

**Synonyms:** CDKN2; p33(CDK2)

Summary: This gene encodes a member of a family of serine/threonine protein kinases that participate

in cell cycle regulation. The encoded protein is the catalytic subunit of the cyclin-dependent protein kinase complex, which regulates progression through the cell cycle. Activity of this protein is especially critical during the G1 to S phase transition. This protein associates with and regulated by other subunits of the complex including cyclin A or E, CDK inhibitor p21Cip1 (CDKN1A), and p27Kip1 (CDKN1B). Alternative splicing results in multiple transcript variants.


[provided by RefSeq, Mar 2014]

**Protein Families:** Druggable Genome, Protein Kinase

Protein Pathways: Cell cycle, Oocyte meiosis, p53 signaling pathway, Pathways in cancer, Progesterone-mediated

oocyte maturation, Prostate cancer, Small cell lung cancer

## **Product images:**



Coomassie blue staining of purified CDK2 protein (Cat# [TP300494]). The protein was produced from HEK293T cells transfected with CDK2 cDNA clone (Cat# [RC200494]) using MegaTran 2.0 (Cat# [TT210002]).