

Product datasheet for TP300495M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CDK1 (NM_001786) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human cell division cycle 2, G1 to S and G2 to M (CDC2), transcript

variant 1, 100 µg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC200495 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MEDYTKIEKIGEGTYGVVYKGRHKTTGQVVAMKKIRLESEEEGVPSTAIREISLLKELRHPNIVSLQDVL MQDSRLYLIFEFLSMDLKKYLDSIPPGQYMDSSLVKSYLYQILQGIVFCHSRRVLHRDLKPQNLLIDDKG TIKLADFGLARAFGIPIRVYTHEVVTLWYRSPEVLLGSARYSTPVDIWSIGTIFAELATKKPLFHGDSEI DQLFRIFRALGTPNNEVWPEVESLQDYKNTFPKWKPGSLASHVKNLDENGLDLLSKMLIYDPAKRISGKM

ALNHPYFNDLDNQIKKM

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 33.9 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 001777

Locus ID: 983



CDK1 (NM_001786) Human Recombinant Protein - TP300495M

UniProt ID: P06493, I6L9I5

RefSeq Size: 1923 Cytogenetics: 10q21.2 891 RefSeq ORF:

Synonyms: CDC2; CDC28A; P34CDC2

Summary: The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This

protein is a catalytic subunit of the highly conserved protein kinase complex known as Mphase promoting factor (MPF), which is essential for G1/S and G2/M phase transitions of eukaryotic cell cycle. Mitotic cyclins stably associate with this protein and function as

regulatory subunits. The kinase activity of this protein is controlled by cyclin accumulation and destruction through the cell cycle. The phosphorylation and dephosphorylation of this protein

also play important regulatory roles in cell cycle control. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar


20091

Protein Families: Druggable Genome, Protein Kinase, Stem cell - Pluripotency

Protein Pathways: Cell cycle, Gap junction, Oocyte meiosis, p53 signaling pathway, Progesterone-mediated

oocyte maturation

Product images:



Coomassie blue staining of purified CDK1 protein (Cat# [TP300495]). The protein was produced from HEK293T cells transfected with CDK1 cDNA clone (Cat# [RC200495]) using MegaTran 2.0

(Cat# [TT210002]).