

Product datasheet for **TP300612**

EIF3B (NM_001037283) Human Recombinant Protein

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant protein of human eukaryotic translation initiation factor 3, subunit B (EIF3B), transcript variant 2, 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC200612 protein sequence Red =Cloning site Green =Tags(s)
	<p>MQDAENVAVPEAAEERAEPGQQQPAAEPPPAEGLLRPAGPGAPEAAGTEASSEEVGIAEAGPEPEVRTEP AAEAEAASGPSESPSPAAEELPGSHAEPVPAQGEAPGEQARDERSDSRAQAVSEDAGGNEGRAAEAEAP RALENGDADEPSFSDPEDFVDDVSEEEELLGDVLKDRPQEAADGIDSVIVVDNVPQVGPDRLEKLNVIHKI FSKFGKITNDFYPEEDGKTKGYIFLEYASPAHAVDAVKNADGYKLDKQHTFRVNLFTDFDKYMTISDEWD IPEKQPFKDLGNLRYWLEEAECRDQYSVIFESGDRTSIFWNDVKDPVSIERARWTETYVVRWSPKGYLA TFHQRGIALWGGGEKFKQIQRFHQGVQLIDFSPCERYLVTFSPMLMDTQDDPQAIWIDILTGHKKRGFHC ESSAHWPIFKWSDGKFFARMTLDTLSIYETPSMGLLDKSLKISGKDFSWSPGGNIIAFVWPEDKDIP ARVTLMQLPTRQEIRVRNLFNVVDCKLHWQKNGDYLCVKVDRTPKGTQGVVTNFEIFRMREKQVPVDVVE MKETIIAFWEPNGSKFAVLHGEAPRISVSFYHVKNNGKIELIKMFDKQANTIFWSPQGQFVLAGLRS MNGALAFVDTSDCTVMNIAEHYMASDVEWDPTGRYVVTSVSWWSHKVDNAYWLWTFQGRLLQKNNKDRFC QLLWRPRPPTLLSQEQIKQIKKDLKKYSKIFEQKDRLSQSKASKELVERRRMMEDFRKYRKMAQELYME QKNERLELRGGVDTDELDSNVDDWEEETIEFFVTEEIIPLGNQE</p> <p>SGPTRTRPLEQKLISEEDLAANDILDYKDDDDKV</p>
Tag:	C-Myc/DDK
Predicted MW:	92.3 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.



[View online »](#)

Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_001032360
Locus ID:	8662
UniProt ID:	P55884 , A0A024R821
RefSeq Size:	3084
Cytogenetics:	7p22.3
RefSeq ORF:	2442
Synonyms:	EIF3-ETA; EIF3-P110; EIF3-P116; EIF3S9; PRT1
Summary:	RNA-binding component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis (PubMed:9388245, PubMed:17581632, PubMed:25849773, PubMed:27462815). The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNA ⁱ and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation (PubMed:9388245, PubMed:17581632). The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression (PubMed:25849773).[UniProtKB/Swiss-Prot Function]

Product images:



Coomassie blue staining of purified EIF3B protein (Cat# TP300612). The protein was produced from HEK293T cells transfected with EIF3B cDNA clone (Cat# [RC200612]) using MegaTran 2.0 (Cat# [TT210002]).