

## **Product datasheet for TP300634**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## BAD (NM 004322) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human BCL2-associated agonist of cell death (BAD), transcript variant

1, 20 μg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC200634 representing NM\_004322

or AA Sequence: Red=Cloning site Green=Tags(s)

MFQIPEFEPSEQEDSSSAERGLGPSPAGDGPSGSGKHHRQAPGLLWDASHQQEQPTSSSHHGGAGAVEI

R

SRHSSYPAGTEDDEGMGEEPSPFRGRSRSAPPNLWAAQRYGRELRRMSDEFVDSFKKGLPRPKSAGTATQ

MRQSSSWTRVFQSWWDRNLGRGSSAPSQ

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

Tag: C-Myc/DDK
Predicted MW: 18.2 kDa

**Concentration:** >0.05 μg/μL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** NP 004313

Locus ID: 572





UniProt ID: Q92934
RefSeq Size: 1127
Cytogenetics: 11q13.1
RefSeq ORF: 504

Synonyms: BBC2; BCL2L8

**Summary:** The protein encoded by this gene is a member of the BCL-2 family. BCL-2 family members are

known to be regulators of programmed cell death. This protein positively regulates cell apoptosis by forming heterodimers with BCL-xL (B-cell lymphoma-extra large) and BCL-2, and reversing their death repressor activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which encode the same

isoform. [provided by RefSeq, Dec 2019]


**Protein Families:** Druggable Genome

**Protein Pathways:** Acute myeloid leukemia, Alzheimer's disease, Amyotrophic lateral sclerosis (ALS), Apoptosis,

Chronic myeloid leukemia, Colorectal cancer, Endometrial cancer, ErbB signaling pathway, Focal adhesion, Insulin signaling pathway, Melanoma, Neurotrophin signaling pathway, Nonsmall cell lung cancer, Pancreatic cancer, Pathways in cancer, Prostate cancer, VEGF signaling

pathway

## **Product images:**



Coomassie blue staining of purified BAD protein (Cat# TP300634). The protein was produced from HEK293T cells transfected with BAD cDNA clone (Cat# [RC200634]) using MegaTran 2.0 (Cat# [TT210002]).