

# Product datasheet for TP300634L

## BAD (NM\_004322) Human Recombinant Protein

### **Product data:**

#### **Product Type: Recombinant Proteins Description:** Recombinant protein of human BCL2-associated agonist of cell death (BAD), transcript variant 1, 1 mg Species: Human **Expression Host:** HEK293T **Expression cDNA Clone** >RC200634 representing NM 004322 or AA Sequence: Red=Cloning site Green=Tags(s) MFQIPEFEPSEQEDSSSAERGLGPSPAGDGPSGSGKHHRQAPGLLWDASHQQEQPTSSSHHGGAGAVEIR SRHSSYPAGTEDDEGMGEEPSPFRGRSRSAPPNLWAAQRYGRELRRMSDEFVDSFKKGLPRPKSAGTATQ MRQSSSWTRVFQSWWDRNLGRGSSAPSQ **TRTRPLEQKLISEEDLAANDILDYKDDDDKV** Tag: C-Myc/DDK Predicted MW: 18.2 kDa **Concentration:** >0.05 µg/µL as determined by microplate BCA method > 80% as determined by SDS-PAGE and Coomassie blue staining Purity: **Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol Recombinant protein was captured through anti-DDK affinity column followed by conventional **Preparation:** chromatography steps. Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process. Store at -80°C. Storage: Stable for 12 months from the date of receipt of the product under proper storage and Stability: handling conditions. Avoid repeated freeze-thaw cycles. **RefSeq:** NP 004313 572 Locus ID: **UniProt ID:** Q92934, A0A024R562



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

	BAD (NM_004322) Human Recombinant Protein – TP300634L
RefSeq Size:	1127
Cytogenetics:	11q13.1
RefSeq ORF:	504
Synonyms:	BBC2; BCL2L8
Summary:	The protein encoded by this gene is a member of the BCL-2 family. BCL-2 family members are known to be regulators of programmed cell death. This protein positively regulates cell apoptosis by forming heterodimers with BCL-xL (B-cell lymphoma-extra large) and BCL-2, and reversing their death repressor activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which encode the same isoform. [provided by RefSeq, Dec 2019]
Protein Families:	Druggable Genome
Protein Pathway	s: Acute myeloid leukemia, Alzheimer's disease, Amyotrophic lateral sclerosis (ALS), Apoptosis, Chronic myeloid leukemia, Colorectal cancer, Endometrial cancer, ErbB signaling pathway, Focal adhesion, Insulin signaling pathway, Melanoma, Neurotrophin signaling pathway, Non- small cell lung cancer, Pancreatic cancer, Pathways in cancer, Prostate cancer, VEGF signaling pathway
Due du et incer	

### **Product images:**

116	_
66	-
45	-
35	-
25	
18	-
14	-

Coomassie blue staining of purified BAD protein (Cat# [TP300634]). The protein was produced from HEK293T cells transfected with BAD cDNA clone (Cat# [RC200634]) using MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US