

Product datasheet for TP300770M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

EIF5A (NM 001970) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human eukaryotic translation initiation factor 5A (EIF5A), transcript

variant B, 100 µg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC200770 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MADDLDFETGDAGASATFPMQCSALRKNGFVVLKGRPCKIVEMSTSKTGKHGHAKVHLVGIDIFTGKKYE DICPSTHNMDVPNIKRNDFQLIGIQDGYLSLLQDSGEVREDLRLPEGDLGKEIEQKYDCGEEILITVLSA

MTEEAAVAIKAMAK

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 16.7 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 001961

Locus ID: 1984

UniProt ID: P63241





RefSeq Size: 1355


Cytogenetics: 17p13.1 RefSeq ORF: 462

Synonyms: eIF-4D; EIF-5A; EIF5A1; eIF5Al

Summary: mRNA-binding protein involved in translation elongation. Has an important function at the

level of mRNA turnover, probably acting downstream of decapping. Involved in actin dynamics and cell cycle progression, mRNA decay and probably in a pathway involved in stress response and maintenance of cell wall integrity. With syntenin SDCBP, functions as a regulator of p53/TP53 and p53/TP53-dependent apoptosis. Regulates also TNF-alphamediated apoptosis. Mediates effects of polyamines on neuronal process extension and survival. May play an important role in brain development and function, and in skeletal muscle stem cell differentiation. Also described as a cellular cofactor of human T-cell leukemia virus type I (HTLV-1) Rex protein and of human immunodeficiency virus type 1 (HIV-1) Rev protein, essential for mRNA export of retroviral transcripts.[UniProtKB/Swiss-Prot Function]

Product images:



Coomassie blue staining of purified EIF5A protein (Cat# [TP300770]). The protein was produced from HEK293T cells transfected with EIF5A cDNA clone (Cat# [RC200770]) using MegaTran 2.0 (Cat# [TT210002]).