

Product datasheet for **TP300866L**

C11orf73 (HIKESHI) (NM_016401) Human Recombinant Protein


Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human chromosome 11 open reading frame 73 (C11orf73), transcript variant 1, 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC200866 protein sequence Red =Cloning site Green =Tags(s) MFGCLVAGRLVQTAAQQAEDKVFVFDLPDYESINHVVVFM LGTIPFPEGMGGSVYFSYPDSNGMPVWQL L GFVTNGKPSAIFKISGLKSGEGSQHPFGAMNIVRTPSVAQIGISVELLD SMAQQTPVGNAAVSSVDSFTQ FTQKMLDNFYNFASSFAVSQAQMTPSPSEMFIPANVVLKWYENFQRR LAQNP LFWKT TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Predicted MW:	21.4 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_057485</u>
Locus ID:	51501


[View online »](#)

UniProt ID:	<u>Q53FT3</u>
RefSeq Size:	1187
Cytogenetics:	11q14.2
RefSeq ORF:	591
Synonyms:	C11orf73; HLD13; HSPC138; HSPC179; L7RN6; OPI10
Summary:	This gene encodes an evolutionarily conserved nuclear transport receptor that mediates heat-shock-induced nuclear import of 70 kDa heat-shock proteins (Hsp70s) through interactions with FG-nucleoporins. The protein mediates transport of the ATP form but not the ADP form of Hsp70 proteins under conditions of heat shock stress. Structural analyses demonstrate that the protein forms an asymmetric homodimer and that the N-terminal domain consists of a jelly-roll/beta-sandwich fold structure that contains hydrophobic pockets involved in FG-nucleoporin recognition. Reduction of RNA expression levels in HeLa cells using small interfering RNAs results in inhibition of heat shock-induced nuclear accumulation of Hsp70s, indicating a role for this gene in regulation of Hsp70 nuclear import during heat shock stress. [provided by RefSeq, Apr 2016]

Product images:



Coomassie blue staining of purified HIKESHI protein (Cat# [TP300866]). The protein was produced from HEK293T cells transfected with HIKESHI cDNA clone (Cat# [RC200866]) using MegaTran 2.0 (Cat# [TT210002]).