

Product datasheet for TP300913

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NEIL2 (NM_145043) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human nei like 2 (E. coli) (NEIL2), transcript variant 1, 20 µg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC200913 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MPEGPLVRKFHHLVSPFVGQQVVKTGGSSKKLQPASLQSLWLQDTQVHGKKLFLRFDLDEEMGPPGSSPT PEPPQKEVQKEGAADPKQVGEPSGQKTLDGSSRSAELVPQGEDDSEYLERDAPAGDAGRWLRVSFGLFGS VWVNDFSRAKKANKRGDWRDPSPRLVLHFGGGGFLAFYNCQLSWSSSPVVTPTCDILSEKFHRGQALEAL GQAQPVCYTLLDQRYFSGLGNIIKNEALYRAGIHPLSLGSVLSASRREVLVDHVVEFSTAWLQGKFQGRP

QHTQVYQKEQCPAGHQVMKEAFGPEDGLQRLTWWCPQCQPQLSEEPEQCQFS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 36.6 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: <u>NP 659480</u> **Locus ID:** 252969



NEIL2 (NM_145043) Human Recombinant Protein - TP300913

UniProt ID: Q969S2, A0A024R361

RefSeq Size: 2746
Cytogenetics: 8p23.1
RefSeq ORF: 996

Synonyms: NEH2; NEI2


Summary: This gene encodes a member of the Fpg/Nei family of DNA glycosylases. These glycosylases

initiate the first step in base excision repair by cleaving oxidatively damaged bases and introducing a DNA strand break via their abasic site lyase activity. This enzyme is primarily associated with DNA repair during transcription and acts prefentially on cytosine-derived lesions, particularly 5-hydroxyuracil and 5-hydroxycytosine. It contains an N-terminal catalytic domain, a hinge region, and a C-terminal DNA-binding domain with helix-two-turn-helix and zinc finger motifs. This enzyme interacts with the X-ray cross complementing factor 1 scaffold protein as part of a multi-protein DNA repair complex. A pseudogene of this gene has been

identified. [provided by RefSeq, Mar 2017]

Protein Families: Druggable Genome
Protein Pathways: Base excision repair

Product images:



Coomassie blue staining of purified NEIL2 protein (Cat# TP300913). The protein was produced from HEK293T cells transfected with NEIL2 cDNA clone (Cat# [RC200913]) using MegaTran 2.0 (Cat# [TT210002]).