

Product datasheet for TP301022

NSFL1C (NM_016143) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human NSFL1 (p97) cofactor (p47) (NSFL1C), transcript variant 1, 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC201022 protein sequence Red=Cloning site Green=Tags(s)

MAAERQEALREFVAVTGAEDRARFFLESAGWDLQIALASFYEDGGDEDIVTISQATPSSVSRGTAPSDN
RVTSFRDLIHDQDEDEEEEEGQRFYAGGSERSGQQIVGPPRKKSPNELVDDLKFGAKEHGAVAVERTKS
PGETSKPRPFAGGGYRLGAAPEEESAYVAGEKRQHSSQDVHVVLKLVKSGFSLDNGELRSYQDPSNAQFL
ESIRRGEVPAELRRLAHGGQVNLDMEDHRDEDFVKPKGAFKAFTGEGQKLGSTAPQVLSTSSPAQQAENE
AKASSILIDESEPTTNIQIRLADGGRLVQKFNHSHRISDIRLFIVDARPAMAATSFILMTTFPNKELAD
ESQTLKEANLLNAVIVQRLT

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	40.4 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_057227</u>



[View online »](#)

Locus ID:	55968
UniProt ID:	Q9UNZ2 , Q53FE8
RefSeq Size:	3568
Cytogenetics:	20p13
RefSeq ORF:	1110
Synonyms:	dj776F14.1; P47; UBX1; UBXD10; UBXN2C
Summary:	N-ethylmaleimide-sensitive factor (NSF) and valosin-containing protein (p97) are two ATPases known to be involved in transport vesicle/target membrane fusion and fusions between membrane compartments. A trimer of the protein encoded by this gene binds a hexamer of cytosolic p97 and is required for p97-mediated regrowth of Golgi cisternae from mitotic Golgi fragments. Alternative splicing results in multiple transcript variants. A related pseudogene has been identified on chromosome 8. [provided by RefSeq, May 2011]

Product images:



Coomassie blue staining of purified NSFL1C protein (Cat# TP301022). The protein was produced from HEK293T cells transfected with NSFL1C cDNA clone (Cat# [RC201022]) using MegaTran 2.0 (Cat# [TT210002]).