

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TP301169

PSMA7 (NM_002792) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins		
Description:	Recombinant protein of human proteasome (prosome, macropain) subunit, alpha type, 7 (PSMA7), 20 μg		
Species:	Human		
Expression Host:	HEK293T		
Expression cDNA Clone or AA Sequence:	>RC201169 protein sequence Red=Cloning site Green=Tags(s)		
	MSYDRAITVFSPDGHLFQVEYAQEAVKKGSTAVGVRGRDIVVLGVEKKSVAKLQDERTVRKICALDDNVC MAFAGLTADARIVINRARVECQSHRLTVEDPVTVEYITRYIASLKQRYTQSNGRRPFGISALIVGFDFDG TPRLYQTDPSGTYHAWKANAIGRGAKSVREFLEKNYTDEAIETDDLTIKLVIKALLEVVQSGGKNIELAV MRRDQSLKILNPEEIEKYVAEIEKEKEENEKKKQKKAS		
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV		
Tag:	C-Myc/DDK		
Predicted MW:	27.7 kDa		
Concentration:	>0.05 μ g/ μ L as determined by microplate BCA method		
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining		
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol		
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.		
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.		
Storage:	Store at -80°C.		
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.		
RefSeq:	<u>NP 002783</u>		
Locus ID:	5688		



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	PSMA7 (NM_002792) Human Recombinant Protein – TP301169
UniProt ID:	<u>014818</u>
RefSeq Size:	1050
Cytogenetics:	20q13.33
RefSeq ORF:	744
Synonyms:	C6; HEL-S-276; HSPC; RC6-1; XAPC7
Summary:	The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. This gene encodes a member of the peptidase T1A family that functions as a 20S core alpha subunit. The encoded protein interacts with the hepatitis B virus X protein and plays a role in regulating hepatitis C virus internal ribosome entry site (IRES) activity, an activity essential for viral replication. The encoded protein also plays a role in the cellular stress response by regulating hypoxia-inducible factor-1alpha. A pseudogene of this gene is located on the long arm of chromosome 9. [provided by RefSeq, Jul 2012]
Protein Families:	Druggable Genome, Protease
Protein Pathways	: Proteasome
Due du et inser	

Product images:

188	_	
98	—	
62	_	
49	-	
38	_	
28	_	-
17	_	
14		
63	=	

Coomassie blue staining of purified PSMA7 protein (Cat# TP301169). The protein was produced from HEK293T cells transfected with PSMA7 cDNA clone (Cat# [RC201169]) using MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US