

Product datasheet for TP301181L

Flavin containing monooxygenase 4 (FMO4) (NM_002022) Human Recombinant Protein

Product data:

| | |
|---------------------------------------|--|
| Product Type: | Recombinant Proteins |
| Description: | Recombinant protein of human flavin containing monooxygenase 4 (FMO4), 1 mg |
| Species: | Human |
| Expression Host: | HEK293T |
| Expression cDNA Clone or AA Sequence: | >RC201181 protein sequence Red =Cloning site Green =Tags(s) |

MAKKVAVIGAGVSGLSSIKCCVDEDLEPTCFERSDDIGGLWKFTESSKDG MTRVYKSLVTNVCKEMSCYS
DFPFHEDYPNFMNHEKFDYDLQFAEHFDLLKYIQFKTTVCSITKRPDFSETGQWDVVTETEGKQNRVAVF
DAVMVCTGHFLNPHLPLEAFPGIHKFKGQILHSQEYKIPEGFQGKRVLVIGLGNTGGDIAVELSRATAAQV
LLSTRGTWVLGRSSDWGYPYNNMMVTRRCCSFIAQVLP SRFLNWIQERKLNKRFNHEDYGLSITKGGKAK
FIVNDELPCILCGAITMKT SVIEFTETSAVFEDGTVEENIDVWIFTTGYTFSFPFFEEPLKSLCTKKIF
LYKQVFPLNLERATLAIIGLIGLKGSI LSGTELQARWVTRVFKGLCKIPPSQKLMMEATEKEQLIKRGVF
KDTSKDKFDYIAYMDDIAACIGTKPSIPL LFLKDPRLAWEVFFGPCTPYQYRLMGP GKWDGARNAILTQW
DRTLKPLKTRIVPDSSKPASMSHYLKAWGAPVLLASLL LICKSSFLKLVRDKLQDRMSPYLVSLWRG

SGPTRTRPLEQKLISEEDLAANDILDYKDDDDKV

| | |
|----------------|--|
| Tag: | C-Myc/DDK |
| Predicted MW: | 63.2 kDa |
| Concentration: | >0.05 µg/µL as determined by microplate BCA method |
| Purity: | > 80% as determined by SDS-PAGE and Coomassie blue staining |
| Buffer: | 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol |
| Preparation: | Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps. |
| Note: | For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process. |
| Storage: | Store at -80°C. |
| Stability: | Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles. |



[View online >](#)

RefSeq: [NP_002013](#)

Locus ID: 2329

UniProt ID: [P31512](#)

RefSeq Size: 2148

Cytogenetics: 1q24.3

RefSeq ORF: 1674


Synonyms: FMO2

Summary: Metabolic N-oxidation of diet-derived amino-trimethylamine (TMA) is mediated by flavin-containing monooxygenase and is subject to an inherited FMO3 polymorphism in man. This results in a small subpopulation with reduced TMA N-oxidation capacity and causes fish odor syndrome (Trimethylaminuria). Three forms of the enzyme are encoded by genes clustered in the 1q23-q25 region. Flavin-containing monooxygenases are NADPH-dependent flavoenzymes that catalyzes the oxidation of soft nucleophilic heteroatom centers in drugs, pesticides, and xenobiotics. [provided by RefSeq, Jan 2015]

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Drug metabolism - cytochrome P450

Product images:



Coomassie blue staining of purified FMO4 protein (Cat# [TP301181]). The protein was produced from HEK293T cells transfected with FMO4 cDNA clone (Cat# [RC201181]) using MegaTran 2.0 (Cat# [TT210002]).