

Product datasheet for TP301200M

Cip4 (TRIP10) (NM_004240) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human thyroid hormone receptor interactor 10 (TRIP10), 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC201200 protein sequence Red=Cloning site Green=Tags(s)

MDWGTTELWDQFEVLERHTQWGLDLLDRYVKFKERTEVEQAYAKQLRSLVKKYLPKRPAKDDPESKFSQQ
QSFVQILQEVNDFAGQRELVAENLSVRVCLELTKYSQEMKQERKMHFQEGRRAQQQLENGFKQLENSKRK
FERDCREAEKAAQTAERLDQDINATKADVEKAKQQAHLRSHMAEESKNEYAAQLQRFNRDQAHFYFSQMP
QIFDKLQDMDERRATRLGAGYGLLSEAELEVPIIAKCLEGMKVAANAVDPKNDSHVLIELHKSGFARPG
DVEFEDFSQPMNRAPSDSSLGTPSDGRPELRGPRSRTRKRWPFGKKNKTVTDFSHLPPEQQRKRLQQQ
LEERSRELQKEVDQREALKKMKDVYEKTPQMGDPASLEPQIAETLSNIERLKLKLEVQKYEAWLAEASRVL
SNRGDSLRSRHARPPDPPASAPPDSSNSASQDTKESSEPPSEESQDTPIYTEFDEDFEEPTSPIGHCV
AIYHFEGSSEGTISMAEGEDLSLMEEDKGDGWTRVRRKEGGEGYVPTSYLRLVTLN

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	62.4 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.



[View online »](#)

RefSeq: [NP_004231](#)

Locus ID: 9322

UniProt ID: [Q15642](#)

RefSeq Size: 2033

Cytogenetics: 19p13.3

RefSeq ORF: 1635


Synonyms: CIP4; HSTP; STOT; STP; TRIP-10

Summary: Required for translocation of GLUT4 to the plasma membrane in response to insulin signaling (By similarity). Required to coordinate membrane tubulation with reorganization of the actin cytoskeleton during endocytosis. Binds to lipids such as phosphatidylinositol 4,5-bisphosphate and phosphatidylserine and promotes membrane invagination and the formation of tubules. Also promotes CDC42-induced actin polymerization by recruiting WASL/N-WASP which in turn activates the Arp2/3 complex. Actin polymerization may promote the fission of membrane tubules to form endocytic vesicles. Required for the formation of podosomes, actin-rich adhesion structures specific to monocyte-derived cells. May be required for the lysosomal retention of FASLG/FASL.[UniProtKB/Swiss-Prot Function]

Protein Families: Druggable Genome

Protein Pathways: Insulin signaling pathway

Product images:



Coomassie blue staining of purified TRIP10 protein (Cat# [TP301200]). The protein was produced from HEK293T cells transfected with TRIP10 cDNA clone (Cat# [RC201200]) using MegaTran 2.0 (Cat# [TT210002]).