

Product datasheet for **TP301248M**

UAP56 (DDX39B) (NM_004640) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human HLA-B associated transcript 1 (BAT1), transcript variant 1, 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC201248 protein sequence Red =Cloning site Green =Tags(s)

MAENDVDNELLDYEDDEVETAAGGDGAEAPAKKDVKGSYVSIHSSGFRDFFLLKPELLRAIVDCGFEHPSE
VQHECIPQAILGMDVLCQAKSGMGKTAVFVLATLQQLEPVTGQVSVLVMCHTRELAFQISKEYERFSKYM
PNVKVAVFFGGLSIKKDEEVLLKNCPHIVVGTPGRILALARNKSLNLKHIFILDECDKMLEQLDMRRD
VQEIFRMTPEHKQVMMFSATLSKEIRPVCVKFMQDPMEIFVDDTKLTLHGLQQYYVKLKDNEKNRKLFD
LLDVLEFNQVWIFVKSVQRCIALAQLLVEQNFPAAIHHRGMPQEERLSRYQQFKDFQRRILVATNLFGRG
MDIERVNI AFNYDMPEDSDTYLHRVARAGRFGTKGLAITFVSDENDAKILNDVQDRFEVNISELPDEIDI
SSYIEQTR

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	48.8 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.



[View online »](#)

RefSeq: [NP_004631](#)

Locus ID: 7919

UniProt ID: [Q13838](#)

RefSeq Size: 2174

Cytogenetics: 6p21.33


RefSeq ORF: 1284

Synonyms: BAT1; D6S81E; UAP56

Summary: This gene encodes a member of the DEAD box family of RNA-dependent ATPases that mediate ATP hydrolysis during pre-mRNA splicing. The encoded protein is an essential splicing factor required for association of U2 small nuclear ribonucleoprotein with pre-mRNA, and it also plays an important role in mRNA export from the nucleus to the cytoplasm. This gene belongs to a cluster of genes localized in the vicinity of the genes encoding tumor necrosis factor alpha and tumor necrosis factor beta. These genes are all within the human major histocompatibility complex class III region. Mutations in this gene may be associated with rheumatoid arthritis. Alternative splicing results in multiple transcript variants. Related pseudogenes have been identified on both chromosomes 6 and 11. Read-through transcription also occurs between this gene and the upstream ATP6V1G2 (ATPase, H⁺ transporting, lysosomal 13kDa, V1 subunit G2) gene. [provided by RefSeq, Feb 2011]

Protein Pathways: Spliceosome

Product images:



Coomassie blue staining of purified DDX39B protein (Cat# [TP301248]). The protein was produced from HEK293T cells transfected with DDX39B cDNA clone (Cat# [RC201248]) using MegaTran 2.0 (Cat# [TT210002]).