

## Product datasheet for TP301259L

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## DRG1 (NM\_004147) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human developmentally regulated GTP binding protein 1 (DRG1), 1

m

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** >RC201259 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MSSTLAKIAEIEAEMARTQKNKATAHHLGLLKARLAKLRRELITPKGGGGGGPGEGFDVAKTGDARIGFV GFPSVGKSTLLSNLAGVYSEVAAYEFTTLTTVPGVIRYKGAKIQLLDLPGIIEGAKDGKGRGRQVIAVAR TCNLILIVLDVLKPLGHKKIIENELEGFGIRLNSKPPNIGFKKKDKGGINLTATCPQSELDAETVKSILA EYKIHNADVTLRSDATADDLIDVVEGNRVYIPCIYVLNKIDQISIEELDIIYKVPHCVPISAHHRWNFDD LLEKIWDYLKLVRIYTKPKGQLPDYTSPVVLPYSRTTVEDFCMKIHKNLIKEFKYALVWGLSVKHNPQKV

**GKDHTLEDEDVIQIVKK** 

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

Tag: C-Myc/DDK
Predicted MW: 40.4 kDa

**Concentration:**  $>0.05 \mu g/\mu L$  as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** NP 004138





Synonyms:

**Locus ID:** 4733

UniProt ID:Q9Y295RefSeq Size:1467Cytogenetics:22q12.2RefSeq ORF:1101


**Summary:** Catalyzes the conversion of GTP to GDP through hydrolysis of the gamma-phosphate bond in

GTP (PubMed:29915238, PubMed:23711155). Appears to have an intrinsic GTPase activity that is stimulated by ZC3H15/DFRP1 binding likely by increasing the affinity for the potassium ions (PubMed:23711155). When hydroxylated at C-3 of 'Lys-22' by JMJD7, may bind to RNA and play a role in translation (PubMed:19819225, PubMed:29915238). Binds to microtubules and promotes microtubule polymerization and stability that are required for mitotic spindle assembly during prophase to anaphase transition. GTPase activity is not necessary for these microtubule-related functions (PubMed:28855639).[UniProtKB/Swiss-Prot Function]

**Protein Families:** Transcription Factors

NEDD3

## **Product images:**



Coomassie blue staining of purified DRG1 protein (Cat# [TP301259]). The protein was produced from HEK293T cells transfected with DRG1 cDNA clone (Cat# [RC201259]) using MegaTran 2.0 (Cat# [TT210002]).