

Product datasheet for **TP301291L**

ABCD4 (NM_005050) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human ATP-binding cassette, sub-family D (ALD), member 4 (ABCD4), transcript variant 1, 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC201291 protein sequence Red =Cloning site Green =Tags(s)

MAVAGPAPGAGARPRLDLQFLQRFLQILKVLFPSSWSSQNALMFLTLLCLLLEQFVIYQVGLIPSQYYGV
LGNKDLEGFKLTFLAVMLIVLNSTLKSFDQFTCNLLYVSWRKDLTEHLHRLYFRGRAYTTLNVLRDDID
NPDQRISQDVERFCRQLSSMASKLIISPFTLVYYTYQCFQSTGWLGPVSIFGYFILGTVVNKTLMGPIVM
KLVHQEKLEGDFRFRKHMQIRVNAEPAAFYRAGHVEHMRDRRLQRLQLTQRELMSKELWLYIGINTFDYL
GSILSYVIAIPIFSGVYGDLSPTLSTLVSKNAFVCIYLISCFTQLIDLSTLSDVAGYTHRIGQLRET
LLDMSLSQDCEILGESKWGLDTPPGWPAAEPADTAFLLERVSISAPSSDKPLIKDLSLKISEGQSLIT
GNTGTGKTSLLRVLGGLWTSTRGSVQMLTDFGPHGVFLPQKPFFTDGLTREQVIYPLKEVYPDSGSADD
ERILRFLELAGLSNLVARTEGLDQQVDWNWYDVLSPGEMQRLSFARLFYLQPKYAVLDEATSALTEEVES
ELYRIGQQLGMTFISVGHRSLEKHFSLVLKLCGGGRWELMRIKVE

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	68.4 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_005041
Locus ID:	5826
UniProt ID:	O14678 , A0A024R6B9
RefSeq Size:	3157
Cytogenetics:	14q24.3
RefSeq ORF:	1818
Synonyms:	ABC41; EST352188; MAHCJ; P70R; P79R; PMP69; PXMP1L
Summary:	The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the ALD subfamily, which is involved in peroxisomal import of fatty acids and/or fatty acyl-CoAs in the organelle. All known peroxisomal ABC transporters are half transporters which require a partner half transporter molecule to form a functional homodimeric or heterodimeric transporter. The function of this peroxisomal membrane protein is unknown. However, it is speculated that it may function as a heterodimer for another peroxisomal ABC transporter and, therefore, may modify the adrenoleukodystrophy phenotype. It may also play a role in the process of peroxisome biogenesis. Alternative splicing results in several protein-coding and non-protein-coding variants. [provided by RefSeq, Jul 2017]
Protein Families:	Druggable Genome, Transmembrane
Protein Pathways:	ABC transporters

Product images:



Coomassie blue staining of purified ABCD4 protein (Cat# [TP301291]). The protein was produced from HEK293T cells transfected with ABCD4 cDNA clone (Cat# [RC201291]) using MegaTran 2.0 (Cat# [TT210002]).