

## Product datasheet for **TP301291M**

### **ABCD4 (NM\_005050) Human Recombinant Protein**

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human ATP-binding cassette, sub-family D (ALD), member 4 (ABCD4), transcript variant 1, 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC201291 protein sequence <b>Red</b> =Cloning site <b>Green</b> =Tags(s)

MAVAGPAPGAGARPRDLQFLQRFLQILKVLFPSSWSSQNALMFLTLLCLLLEQFVIYQVGLIPSQYYGV  
LGNKDLEGFKTLTFLAVMLIVLNSTLKSFDQFTCNLLYVSWRKDLTEHLHRLYFRGRAYTTLNVLRDDID  
NPDQRISQDVERFCRQLSSMASKLIISPFTLVYYTYQCFQSTGWLGPVSIFGYFILGTVVNKTLMGPIVM  
KLVHQEKLEGDFRFKHMQIRVNAEPAAFYRAGHVEHMRDRRLQRLQLTQRELMSKELWLYIGINTFDYL  
GSILSYVIAIPIFSGVYGDLSPTLSTLVSKNAFVCIYLISCFTQLIDLSTLSDVAGYTHRIGQLRET  
LLDMSLSQDCEILGESKWGLDTPPGWPAEAPADTAFLLERVSISAPSSDKPLIKDLSLKISEGQSLIT  
GNTGTGKTSLLRVLGGLWTSTRGSVQMLTDFGPHGVFLPQKPFFTDGLTREQVIYPLKEVYPDSGSADD  
ERILRFLELAGLSNLVARTEGLDQQVDWNWYDVLSPGEMQRLSFARLFYLQPKYAVLDEATSALTEEVES  
ELYRIGQQLGMTFISVGHRSLEKHFSLVLKLCGGGRWELMRIKVE

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV**


Tag:	C-Myc/DDK
Predicted MW:	68.4 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

<b>Stability:</b>	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
<b>RefSeq:</b>	<a href="#">NP_005041</a>
<b>Locus ID:</b>	5826
<b>UniProt ID:</b>	<a href="#">O14678</a> , <a href="#">A0A024R6B9</a>
<b>RefSeq Size:</b>	3157
<b>Cytogenetics:</b>	14q24.3
<b>RefSeq ORF:</b>	1818
<b>Synonyms:</b>	ABC41; EST352188; MAHCJ; P70R; P79R; PMP69; PXMP1L
<b>Summary:</b>	The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the ALD subfamily, which is involved in peroxisomal import of fatty acids and/or fatty acyl-CoAs in the organelle. All known peroxisomal ABC transporters are half transporters which require a partner half transporter molecule to form a functional homodimeric or heterodimeric transporter. The function of this peroxisomal membrane protein is unknown. However, it is speculated that it may function as a heterodimer for another peroxisomal ABC transporter and, therefore, may modify the adrenoleukodystrophy phenotype. It may also play a role in the process of peroxisome biogenesis. Alternative splicing results in several protein-coding and non-protein-coding variants. [provided by RefSeq, Jul 2017]
<b>Protein Families:</b>	Druggable Genome, Transmembrane
<b>Protein Pathways:</b>	ABC transporters

### Product images:



Coomassie blue staining of purified ABCD4 protein (Cat# [TP301291]). The protein was produced from HEK293T cells transfected with ABCD4 cDNA clone (Cat# [RC201291]) using MegaTran 2.0 (Cat# [TT210002]).