

Product datasheet for **TP301314M**

Bcl x (BCL2L1) (NM_138578) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human BCL2-like 1 (BCL2L1), nuclear gene encoding mitochondrial protein, transcript variant 1, 100 µg

Species: Human

Expression Host: HEK293T

Expression cDNA Clone or AA Sequence: >RC201314 protein sequence
Red=Cloning site **Green**=Tags(s)

MSQSNRELWVDFLSYKLSQKGYWSQFSDVEENRTEAPEGTESEMETPSAINGNPSWHLADSPAVNGATG
HSSSLDAREVIPMAAVKQALREAGDEFELRYRRAFSDLTSQLHITPGTAYQSFEQVVNELFRDGVNWGRI
VAFFSFGGALCVESVDKEMQVLVSRIAAMATYLNHLEPWIQENGGWDTFVELYGNNAAESRKGQERF
NRWFLTGMTVAGVVLLGSLFSRK

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 25.9 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Bioactivity: In vitro ubiquitination assay substrate (PMID: [28038320](#))

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: [NP_612815](#)



[View online »](#)


Locus ID: 598
UniProt ID: [Q07817](#), [Q07817-1](#), [A0A0S2Z3C5](#)
RefSeq Size: 2575
Cytogenetics: 20q11.21
RefSeq ORF: 699
Synonyms: Bcl-X; BCL-XL/S; BCL2L; BCLX; PPP1R52

Summary: The protein encoded by this gene belongs to the BCL-2 protein family. BCL-2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. The proteins encoded by this gene are located at the outer mitochondrial membrane, and have been shown to regulate outer mitochondrial membrane channel (VDAC) opening. VDAC regulates mitochondrial membrane potential, and thus controls the production of reactive oxygen species and release of cytochrome C by mitochondria, both of which are the potent inducers of cell apoptosis. Alternative splicing results in multiple transcript variants encoding two different isoforms. The longer isoform acts as an apoptotic inhibitor and the shorter isoform acts as an apoptotic activator. [provided by RefSeq, Dec 2015]


Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Transmembrane

Protein Pathways: Amyotrophic lateral sclerosis (ALS), Apoptosis, Chronic myeloid leukemia, Jak-STAT signaling pathway, Pancreatic cancer, Pathways in cancer, Small cell lung cancer

Product images:



PARK2 E3 ubiquitin ligase ubiquitinates BCL-XL in vitro. Recombinant proteins BCL-XL (OriGene [TP301314]) and PARK2 were incubated in HEPES buffer, pH 8.0 containing E1, E2, ubiquitin, and Mg²⁺-ATP. Reactions were cultured for 10 min at 30 C, then 30 - 60 min at 37 C, and reaction samples were subjected to Western blot analysis. Figure cited from Neoplasia, PMID: 28038320



Coomassie blue staining of purified BCL2L1 protein (Cat# [TP301314]). The protein was produced from HEK293T cells transfected with BCL2L1 cDNA clone (Cat# [RC201314]) using MegaTran 2.0 (Cat# [TT210002]).